

Product datasheet for SC316362

CDK10 (NM_001098533) Human Untagged Clone

Product data:

OriGene Technologies, Inc.


9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Expression Plasmids
Product Name:	CDK10 (NM_001098533) Human Untagged Clone
Tag:	Tag Free
Symbol:	CDK10
Synonyms:	ALSAS; PISSLRE
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC316362 representing NM_001098533. Blue=Insert sequence <mark>Red</mark> =Cloning site Green=Tag(s)
	GCTCGTTTAGTGAACCGTCAGAATTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTG GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC ATGGACAAGGAGAAGGATGGCATCCCCATCAGCAGCTTGCGGGGAGATCACGCTGCTGCTGCCGCCTGCGT CATCCGAACATCGTGGAGCTGAAGGAGGTGGTTGTGGGGGAACCACCTGGAGAGCATCTTCCTGGTGATG GGTTACTGTGAGCAGGACCTGGCCAGCCTCCTGGAGAATATGCCAACACCCTTCTCGGAGGGCTCAGGTC AAGTGCATCGTGCTGCAGGTGCTCCGGGGCCTCCAGTATCTGCACAGGAACTTCATTATCCACAGGGAC CTGAAGGTTTCCAACTTGCTCATGACCGACAAGGGTTGTGTGAAGACAGCGGATTTCGGCCTGGCCCGG GCCTATGGTGTCCCAGTAAAGCCAATGACCCCCAAGGTGGTCACTCTCTGGTACCGAGCCCCTGAACTG CTGTTGGGAACCACCACGCAGACCACCAGCATCGACATGTGGGGCTGCGGCTGCATACTGGCCGAGCTG GGCACGCCCAGTGAGAACATCTGGCCGGGCTTTTCCAAGGTGGCCGGCTGCGACACTGGCCGGGC AAGCAGCCTACAACAACCTGAAGCACAAGTTCCCATGGCTGCGGCAGCAGGCCGGCTGCGCCAGCTGCG GGCACGCCCAGTGAGAACATCTGGCCGGGGCTTTTCCCAAGCTGCCGGCGGCTGCGCCAGTACAGCCTCCGG AAGCAGCCCTACAACAACCTGAAGCACAAGTTCCCATGGCTGTCGGAGGCCGGCC
Restriction Sites:	Sgfl-Mlul



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Plasmid Map:



ACCN:	NM_001098533
Insert Size:	852 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	 Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM 001098533.2</u>
RefSeq Size:	1759 bp
RefSeq ORF:	852 bp

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	CDK10 (NM_001098533) Human Untagged Clone – SC316362
Locus ID:	8558
UniProt ID:	<u>Q15131</u>
Cytogenetics:	16q24.3
Protein Families	: Druggable Genome, Protein Kinase
MW:	31.9 kDa
Gene Summary:	The protein encoded by this gene belongs to the CDK subfamily of the Ser/Thr protein kinase family. The CDK subfamily members are highly similar to the gene products of S. cerevisiae cdc28, and S. pombe cdc2, and are known to be essential for cell cycle progression. This kinase has been shown to play a role in cellular proliferation and its function is limited to cell cycle G2-M phase. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, May 2009] Transcript Variant: This variant (c) uses an alternate donor splice site at the first exon resulting in translation from a downstream start codon, and uses an alternate donor splice site in the 3' coding region, compared to variant a. The resulting isoform (c) has a shorter N-terminus and lacks a 6-aa segment near the C-terminus, compared to isoform a.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US