

Product datasheet for SC316703

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

ICAM2 (NM_001099788) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: ICAM2 (NM_001099788) Human Untagged Clone

Tag: Tag Free
Symbol: ICAM2
Synonyms: CD102

Vector: pCMV6 series

Fully Sequenced ORF: >NCBI ORF sequence for NM_001099788, the custom clone sequence may differ by one or

more nucleotides

GGGGTGCGAGCGCTTGGAGGAGGCTGCCCAGGCCTTCCGGCCA

Restriction Sites: Please inquire **ACCN:** NM 001099788

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001099788.1</u>, <u>NP 001093258.1</u>

 RefSeq Size:
 1235 bp

 RefSeq ORF:
 828 bp

 Locus ID:
 3384

 UniProt ID:
 P13598

 Cytogenetics:
 17q23.3

Protein Families: ES Cell Differentiation/IPS, Transmembrane

Protein Pathways: Cell adhesion molecules (CAMs), Natural killer cell mediated cytotoxicity

Gene Summary: The protein encoded by this gene is a member of the intercellular adhesion molecule (ICAM)

family. All ICAM proteins are type I transmembrane glycoproteins, contain 2-9

immunoglobulin-like C2-type domains, and bind to the leukocyte adhesion LFA-1 protein. This

protein may play a role in lymphocyte recirculation by blocking LFA-1-dependent cell

adhesion. It mediates adhesive interactions important for antigen-specific immune response,

NK-cell mediated clearance, lymphocyte recirculation, and other cellular interactions important for immune response and surveillance. Several transcript variants encoding the

same protein have been found for this gene. [provided by RefSeq, Jul 2008]

Transcript Variant: This variant (3) differs in the 5' UTR compared to variant 1. All 5 variants

encode the same protein.