

Product datasheet for SC316909

ZFAND5 (NM 001102420) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: ZFAND5 (NM 001102420) Human Untagged Clone

Tag: Tag Free Symbol: ZFAND5

Synonyms: ZA20D2; ZFAND5A; ZNF216

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >SC316909 representing NM_001102420.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT

TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

Restriction Sites: Sgfl-Mlul

Plasmid Map:

ACCN: NM_001102420

Insert Size: 642 bp



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



ZFAND5 (NM_001102420) Human Untagged Clone - SC316909

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001102420.2</u>

 RefSeq Size:
 5887 bp

 RefSeq ORF:
 642 bp

 Locus ID:
 7763

 UniProt ID:
 076080

Cytogenetics: 9q21.13

MW: 23.1 kDa

Gene Summary: Involved in protein degradation via the ubiquitin-proteasome system. May act by anchoring

ubiquitinated proteins to the proteasome. Plays a role in ubiquitin-mediated protein degradation during muscle atrophy. Plays a role in the regulation of NF-kappa-B activation and apoptosis. Inhibits NF-kappa-B activation triggered by overexpression of RIPK1 and TRAF6 but not of RELA. Inhibits also tumor necrosis factor (TNF), IL-1 and TLR4-induced NF-kappa-B activation in a dose-dependent manner. Overexpression sensitizes cells to TNF-induced apoptosis. Is a potent inhibitory factor for osteoclast differentiation.[UniProtKB/Swiss-Prot

Function]

Transcript Variant: This variant (a) represents the longest transcript. All six variants encode the same protein. Sequence Note: This RefSeq record was created from transcript and genomic sequence data because no single transcript was available for the full length of the

gene. The extent of this transcript is supported by transcript alignments.