

## Product datasheet for **SC317007**

### **ECHDC1 (NM\_001105545) Human Untagged Clone**

#### **Product data:**

<b>Product Type:</b>	Expression Plasmids
<b>Product Name:</b>	ECHDC1 (NM_001105545) Human Untagged Clone
<b>Tag:</b>	Tag Free
<b>Symbol:</b>	ECHDC1
<b>Synonyms:</b>	dj351K20.2; HEL-S-76; MMCD
<b>Vector:</b>	<u>pCMV6 series</u>
<b>Fully Sequenced ORF:</b>	>NCBI ORF sequence for NM_001105545, the custom clone sequence may differ by one or more nucleotides ATGATGCTACAACCTTCTGGAAAAAGTAATTGAATTGAAAATTGGACAGAGGGGAAAGGC CTCATTGTCCGTGGGGCAAAAAATACTTTCTCTTCAGGATCTGATCTGAATGCTGTGAAA TCACTAGGAACTCCAGAGACTTCCTTTAATAAGTGTTCGCTGGTTCAAGGTTGGGCATT GGGTGGAGGAGCAGAATTTACTACAGCATG
<b>Restriction Sites:</b>	Please inquire
<b>ACCN:</b>	NM_001105545
<b>OTI Disclaimer:</b>	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
<b>OTI Annotation:</b>	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>



[View online »](#)

RefSeq: [NM\\_001105545.1](#), [NP\\_001099015.1](#)

RefSeq Size: 2079 bp

RefSeq ORF: 213 bp

Locus ID: 55862

UniProt ID: [Q9NTX5](#)

Cytogenetics: 6q22.33

**Gene Summary:** Decarboxylases ethylmalonyl-CoA decarboxylase, a potentially toxic metabolite, to form butyryl-CoA, suggesting it might be involved in metabolite proofreading. Also has methylmalonyl-CoA decarboxylase activity at lower level.[UniProtKB/Swiss-Prot Function]  
Transcript Variant: This variant (4) lacks two alternate exons compared to variant 1, which causes a frameshift and a downstream AUG start codon to be used. The resulting isoform (4) has a distinct C-terminus and is shorter at both the N- and C-termini compared to isoform 1.