

Product datasheet for SC317047

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GJB6 (NM_001110220) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: GJB6 (NM_001110220) Human Untagged Clone

Tag: Tag Free Symbol: G|B6

Synonyms: CX30; DFNA3; DFNA3B; DFNB1B; ECTD2; ED2; EDH; HED; HED2

Vector: <u>pCMV6 series</u>

Fully Sequenced ORF: >NCBI ORF sequence for NM_001110220, the custom clone sequence may differ by one or

more nucleotides

AGC

Restriction Sites: Please inquire **ACCN:** NM 001110220

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeg: NM 001110220.1, NP 001103690.1

 RefSeq Size:
 1990 bp

 RefSeq ORF:
 786 bp

 Locus ID:
 10804

 UniProt ID:
 095452

 Cytogenetics:
 13q12.11

Protein Families: Druggable Genome, Transmembrane

Gene Summary: Gap junctions allow the transport of ions and metabolites between the cytoplasm of adjacent

cells. They are formed by two hemichannels, made up of six connexin proteins assembled in groups. Each connexin protein has four transmembrane segments, two extracellular loops, a cytoplasmic loop formed between the two inner transmembrane segments, and the N- and C-terminus both being in the cytoplasm. The specificity of the gap junction is determined by which connexin proteins comprise the hemichannel. In the past, connexin protein names were based on their molecular weight, however the new nomenclature uses sequential numbers based on which form (alpha or beta) of the gap junction is present. This gene encodes one of the connexin proteins. Mutations in this gene have been found in some forms

2008]

Transcript Variant: This variant (2) lacks a segment in the 5' UTR, compared to variant 1. All

of deafness and in some families with hidrotic ectodermal dysplasia. [provided by RefSeq, Jul

variants encode the same protein.