

Product datasheet for SC317926

MAPK15 (NM_139021) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: MAPK15 (NM_139021) Human Untagged Clone

Tag: Tag Free Symbol: MAPK15

Synonyms: ERK7; ERK8

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn





Fully Sequenced ORF:

>SC317926 representing NM_139021.

Blue=Insert sequence Red=Cloning site Green=Tag(s)


ATGTGCACCGTAGTGGACCCTCGCATTGTCCGGAGATACCTACTCAGGCGGCAGCTCGGGCAGGGGGGCC TATGGCATTGTGTGGAAGGCAGTGGACCGGAGGACTGGTGAGGTCGTGGCCATCAAGAAAATCTTTGAT GCTTTTAGGGATAAGACAGATGCCCAGAGAACATTCCGGGAAATCACGCTCCTCCAGGAGTTTGGGGAC CATCCCAACATCATCAGCCTCCTTGACGTGATCCGGGCAGAGAACGACAGGGACATTTACCTGGTGTTT GAGTTTATGGACACTGACCTGAACGCAGTCATCCGGAAGGGCGGCCTGCTGCAGGACGTCCACGTGCGC TCCATCTTCTACCAGCTCCTGCGGGCCACCCGGTTCCTCCACTCGGGGCACGTTGTGCACCGGGACCAG AAGCCGTCCAATGTGCTCCTGGATGCCAACTGCACAGTGAAGCTGTGTGACTTTGGCCTGGCCCGCTCC CTGGGCGACCTCCCCGAGGGGCCTGAGGACCAGGCCGTGACAGAGTACGTGGCCACACGCTGGTACCGA GCACCGGAGGTGCTGCTCTCTCGCACCGATACACCCTTGGGGTGGACATGTGGAGTCTGGGCTGTATC CTGGGGGAGATGCTGCGGGGGAGACCCCTGTTCCCCGGCACGTCCACCCTCCACCAGCTGGAGCTGATC CTGGAGACCATCCCACCGCCATCTGAGGAGGACCTCCTGGCTCTCGGCTCAGGCTGCCGTGCCTCTGTG CTGCACCAGCTGGGGTCCCGGCCACGACAGACGCTGGATGCCCTCCTACCGCCAGACACCTCCCCAGAG GCCTTGGACCTCCTTAGGCGACTCCTGGTGTTCGCCCCGGACAAGCGGTTAAGCGCGACCCAGGCACTG CAGCACCCCTACGTGCAGAGGTTCCACTGCCCCAGCGACGAGTGGGCACGAGAGGCAGATGTGCGGCCC CGGGCACACGAAGGGGTCCAGCTCTCTGTGCCTGAGTACCGCAGCCGCGTCTATCAGATGATCCTGGAG TGTGGAGGCAGCAGCGGCACCTCGAGAGAGAGGGCCCGGAGGGTGTCTCCCCAAGCCAGGCACACCTG CACAAACCCAGAGCCGACCCTCAGCTGCCTTCTAGGACACCTGTGCAGGGTCCCAGACCCAGGCCCCAG AGCAGCCCAGGCCATGACCCTGCCGAGCACGAGTCCCCCCGTGCAGCCAAGAACGTTCCCAGGCAGAAC GCTCAGGTGGCCAACCAGGCCCTGATCCGGGGTGACTGGAACCGGGGCGGTGGGGTGAGGGTGGCCAGC GTACAACAGGTCCCTCCCGGCTTCCTCCGGAGGCCCGGCCCGGAGGATGTTCAGCACCTCTGCC TTGCAGGGTGCCCAGGGGTGCCAGGGCTTTGCTTGGAGGCTACTCCCAAGCCTACGGGACTGTCTGC CACTCGGCACTGGCCACCTGCCCCTGCTGGAGGGGCACCATGTGTGA

ACGCGTACGCCGCCCCCCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

Restriction Sites:

Plasmid Map:

Sgfl-Mlul



ACCN: NM_139021 **Insert Size:** 1635 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 139021.2</u>

 RefSeq Size:
 1904 bp

 RefSeq ORF:
 1635 bp

 Locus ID:
 225689

 UniProt ID:
 Q8TD08

 Cytogenetics:
 8q24.3

Protein Families: Druggable Genome, Protein Kinase

MW: 59.8 kDa



Gene Summary:

Atypical MAPK protein that regulates several process such as autophagy, ciliogenesis, protein trafficking/secretion and genome integrity, in a kinase activity-dependent manner (PubMed:22948227, PubMed:24618899, PubMed:29021280, PubMed:21847093, PubMed:20733054). Controls both, basal and starvation-induced autophagy throught its interaction with GABARAP, MAP1LC3B and GABARAPL1 leading to autophagosome formation, SQSTM1 degradation and reduced MAP1LC3B inhibitory phosphorylation (PubMed:22948227). Regulates primary cilium formation and the localization of ciliary proteins involved in cilium structure, transport, and signaling (PubMed:29021280). Prevents the relocation of the sugar-adding enzymes from the Golgi to the endoplasmic reticulum, thereby restricting the production of sugar-coated proteins (PubMed:24618899). Upon aminoacid starvation, mediates transitional endoplasmic reticulum site disassembly and inhibition of secretion (PubMed:21847093). Binds to chromatin leading to MAPK15 activation and interaction with PCNA, that which protects genomic integrity by inhibiting MDM2-mediated degradation of PCNA (PubMed:20733054). Regulates DA transporter (DAT) activity and protein expression via activation of RhoA (PubMed:28842414). In response to H(2)O(2) treatment phosphorylates ELAVL1, thus preventing it from binding to the PDCD4 3' UTR and rendering the PDCD4 mRNA accessible to miR-21 and leading to its degradation and loss of protein expression (PubMed:26595526). Also functions in a kinase activity-independent manner as a negative regulator of growth (By similarity). Phosphorylates in vitro FOS and MBP (PubMed:11875070, PubMed:16484222, PubMed:20638370, PubMed:19166846). During oocyte maturation, plays a key role in the microtubule organization and meiotic cell cycle progression in oocytes, fertilized eggs, and early embryos (By similarity). Interacts with ESRRA promoting its re-localization from the nucleus to the cytoplasm and then prevents its transcriptional activity (PubMed:21190936).[UniProtKB/Swiss-Prot Function]