

## **Product datasheet for SC318771**

## IGF1 (NM 001111284) Human Untagged Clone

## **Product data:**

**Product Type:** Expression Plasmids

Product Name: IGF1 (NM\_001111284) Human Untagged Clone

Tag: Tag Free Symbol: IGF1

Synonyms: IGF; IGF-I; IGFI; MGF

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >SC318771 representing NM\_001111284.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

**ACGCGTACGCGCCCCTC**GAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT

TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

Restriction Sites: Sgfl-Mlul

ACCN: NM 001111284

**Insert Size:** 414 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



**OTI Annotation:** 

This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001111284.1</u>

 RefSeq Size:
 7204 bp

 RefSeq ORF:
 414 bp

 Locus ID:
 3479

 UniProt ID:
 P05019

 Cytogenetics:
 12q23.2

**Protein Families:** Druggable Genome, ES Cell Differentiation/IPS, Secreted Protein

Protein Pathways: Dilated cardiomyopathy, Focal adhesion, Glioma, Hypertrophic cardiomyopathy (HCM), Long-

term depression, Melanoma, mTOR signaling pathway, Oocyte meiosis, p53 signaling pathway, Pathways in cancer, Progesterone-mediated oocyte maturation, Prostate cancer

**MW:** 15.2 kDa

**Gene Summary:** The protein encoded by this gene is similar to insulin in function and structure and is a

member of a family of proteins involved in mediating growth and development. The encoded protein is processed from a precursor, bound by a specific receptor, and secreted. Defects in this gene are a cause of insulin-like growth factor I deficiency. Alternative splicing results in multiple transcript variants encoding different isoforms that may undergo similar processing

to generate mature protein. [provided by RefSeq, Sep 2015]

Transcript Variant: This variant (2) differs in the 5' UTR and has multiple coding region differences, compared to variant 1. These differences cause translation initiation from a distinct start codon and result in an isoform (2) with novel N- and C-termini, compared to isoform 1. This isoform is also known as IIA. This isoform (2) may undergo proteolytic processing similar to isoform 1. Sequence Note: This RefSeq was created from transcript and genomic sequence because transcript sequence consistent with the reference assembly was not available for all regions of the RefSeq transcript. The extent of this transcript is supported by transcript alignments.

by transcript alignments.