

Product datasheet for **SC318776**

HIP2 (UBE2K) (NM_001111113) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	HIP2 (UBE2K) (NM_001111113) Human Untagged Clone
Tag:	Tag Free
Symbol:	UBE2K
Synonyms:	E2-25K; HIP2; HYPG; LIG; UBC1
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC318776 representing NM_001111113. Blue=Insert sequence Red=Cloning site Green=Tag(s)


```
GCTCGTTTAGTGAACCGTCAGAATTTGTAAACGACTCACTATAGGGCGCCGGGAATTCGTCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCCGATCGCC
ATGGCCAACATCGCGGTGCAGCGAATCAAGCGGGAGTTCAAGGAGGTGCTGAAGAGCGAGGAGTCCGG
TTTATCACTAAAATATGGCATCCTAATATTAGTTCGTCACAGGGGCTATTTGTTGGATATCCTGAAA
GATCAATGGGCAGCTGCAATGACTCTCCGACGGTATTATTGTCATTGCAAGCACTATTGGCAGCTGCA
GAGCCAGATGATCCACAGGATGCTGTAGTAGCAAATCAGTACAAACAAAATCCCGAAATGTTCAAACAG
ACAGCTCGACTTTGGGCACATGTGTATGCTGGAGCACCAGTTTCTAGTCCAGAATACACCAAAAAATA
GAAAACCTATGTGCTATGGGCTTTGATAGGAATGCAGTAATAGTGCCTTGCTCTCAAAATCATGGGAT
GTAGAGACTGCAACAGAATTGCTTCTGAGTAACTGA
ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
```

Restriction Sites: Sgfl-Mlul



[View online »](#)

Plasmid Map:



ACCN:	NM_001111113
Insert Size:	450 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001111113.1](#)

RefSeq Size: 5106 bp

RefSeq ORF: 450 bp

Locus ID: 3093

UniProt ID: [P61086](#)

Cytogenetics: 4p14

Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: Ubiquitin mediated proteolysis

MW: 16.6 kDa

Gene Summary: The protein encoded by this gene belongs to the ubiquitin-conjugating enzyme family. This protein interacts with RING finger proteins, and it can ubiquitinate huntingtin, the gene product for Huntington's disease. Known functions for this protein include a role in aggregate formation of expanded polyglutamine proteins and the suppression of apoptosis in polyglutamine diseases, a role in the dislocation of newly synthesized MHC class I heavy chains from the endoplasmic reticulum, and involvement in foam cell formation. Multiple transcript variants encoding different isoforms have been identified for this gene. [provided by RefSeq, Jul 2008]

Transcript Variant: This variant (3) lacks an alternate in-frame segment in the 5' coding region, compared to variant 1, resulting in a shorter protein (isoform 3), compared to isoform 1.

Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.