

Product datasheet for SC318799

NACA (NM_001113201) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	NACA (NM_001113201) Human Untagged Clone
Tag:	Tag Free
Symbol:	NACA
Synonyms:	HSD48; NAC-alpha; NACA1; skNAC
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC318799 representing NM_001113201. Blue=Insert sequence Red=Cloning site Green=Tag(s)

GCTCGTTTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTG
 GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
 ATGCCCGGCGAAGCCACAGAAACCGTCCCTGCTACAGAGCAGGAGTTGCCGCAGCCCCAGGCTGAGACA
 GGGTCTGGAACAGAATCTGACAGTGATGAATCAGTACCAGAGCTTGAAGAACAGGATTCCACCCAGGCA
 ACCACACAACAAGCCCAGCTGGCGGCAGCAGCTGAAATTGATGAAGAACCAGTCAGTAAAGCAAAACAG
 AGTCGGAGTGAAGAAGGCACGGAAGGCTATGTCCAACTGGGTCTTCGGCAGGTTACAGGAGTTACT
 AGAGTCACTATCCGGAATCTAAGAATATCCTCTTTGTCATCACAAAACCAGATGTCTACAAGAGCCCT
 GCTTCAGATACTTACATAGTTTTTGGGAAGCCAAGATCGAAGATTTATCCAGCAAGCACAACTAGCA
 GCTGCTGAGAAATTCAAAGTTCAAGGTGAAGCTGTCTCAACATTCAAGAAAACACAGACTCCAAC
 GTACAAGAGGAGAGTGAAGAGGAAGAGGTCGATGAAACAGGTGTAGAAGTTAAGGACATTGAATTGGTC
 ATGTCACAAGCAAATGTGTCGAGAGCAAAGGCAGTCCGAGCCCTGAAGAACAACAGTAATGATATTGTA
 AATGCGATTATGGAATTAACAATGTAA
 ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
 TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

Restriction Sites:	SgfI-MluI
ACCN:	NM_001113201
Insert Size:	648 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).


[View online »](#)

OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001113201.2
RefSeq Size:	1138 bp
RefSeq ORF:	648 bp
Locus ID:	4666
UniProt ID:	Q13765
Cytogenetics:	12q13.3
Protein Families:	Druggable Genome, Transcription Factors
MW:	23.4 kDa
Gene Summary:	<p>This gene encodes a protein that associates with basic transcription factor 3 (BTF3) to form the nascent polypeptide-associated complex (NAC). This complex binds to nascent proteins that lack a signal peptide motif as they emerge from the ribosome, blocking interaction with the signal recognition particle (SRP) and preventing mistranslocation to the endoplasmic reticulum. This protein is an IgE autoantigen in atopic dermatitis patients. Alternative splicing results in multiple transcript variants, but the full length nature of some of these variants, including those encoding very large proteins, has not been determined. There are multiple pseudogenes of this gene on different chromosomes. [provided by RefSeq, Feb 2016]</p> <p>Transcript Variant: This variant (2) contains multiple differences in the 5' UTR and lacks three alternate exons in the central coding region, but maintains the reading frame, compared to variant 1. This variant encodes isoform b, which is shorter than isoform a. Variants 2, 3, 4, and 5, encode the same isoform (b).</p>