

Product datasheet for SC318814

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MTRF1L (NM_001114184) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: MTRF1L (NM 001114184) Human Untagged Clone

Tag: Tag Free
Symbol: MTRF1L

Synonyms: HMRF1L; MRF1L; mtRF1a

Vector: <u>pCMV6 series</u>

Fully Sequenced ORF: >NCBI ORF sequence for NM_001114184, the custom clone sequence may differ by one or

more nucleotides

GTCCGGATAGTTCATCTTCCAACAGATTGGAAG

Restriction Sites: Please inquire ACCN: NM 001114184

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).







Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeg: NM 001114184.1, NP 001107656.1

RefSeq Size:3741 bpRefSeq ORF:816 bpLocus ID:54516UniProt ID:Q9UGC7Cytogenetics:6q25.2

Gene Summary: The protein encoded by this gene plays a role in mitochondrial translation termination, and is

thought to be a release factor that is involved in the dissociation of the complete protein from the final tRNA, the ribosome, and the cognate mRNA. This protein acts upon UAA and UAG stop codons, but has no in vitro activity against UGA, which encodes tryptophan in human mitochondrion, or, the mitochondrial non-cognate stop codons, AGA and AGG. This protein shares sequence similarity to bacterial release factors. Pseudogenes of this gene are found on chromosomes 4, 8, and 11. Alternative splicing results in multiple transcript

variants. [provided by RefSeq, Sep 2014]

Transcript Variant: This variant (3) lacks an alternate exon in the 3' coding region, which results in a frameshift and an early stop codon, compared to variant 1. This variant encodes

isoform 3, which is shorter and has a distinct C-terminus, compared to isoform 1.