

Product datasheet for **SC319433**

Apolipoprotein E (APOE) (NM_000041) Human Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: Apolipoprotein E (APOE) (NM_000041) Human Untagged Clone
Tag: Tag Free
Symbol: Apolipoprotein E
Synonyms: AD2; APO-E; ApoE4; LDLCQ5; LPG
Mammalian Cell Selection: Neomycin
Vector: pCMV6-AC (PS100020)
E. coli Selection: Ampicillin (100 ug/mL)

Fully Sequenced ORF: >OriGene sequence for NM_000041.2
GGACGTCCTTCCCAGGAGCCGACTGGCCAATCACAGGCAGGAAGATGAAGTTCTGTGG
GCTGCGTTGCTGGTCACATTCTGGCAGGATGCCAGGCCAAGGTGGAGCAAGCGTGGAG
ACAGAGCCGGAGCCCGAGCTGCGCCAGCAGACCGAGTGGCAGAGCGGCCAGCGCTGGGAA
CTGGCACTGGGTGCTTTTGGGATTACCTGCGCTGGGTGCAGACTGTCTGAGCAGGTG
CAGGAGGAGCTGCTCAGCTCCCAGGTACCCAGGAACTGAGGGCGCTGATGGACGAGACC
ATGAAGGAGTTGAAGGCCTACAAATCGGAACTGGAGGAACAACCTGACCCCGTGGCGGAG
GAGACGCGGGCACGGCTGTCCAAGGAGCTGCAGGCGGCGCAGGCCCGGCTGGGCGCGGAC
ATGGAGGACGTGTGCGGCCGCTGGTGCAGTACCGCGGCGAGGTGCAGGCCATGCTCGGC
CAGAGCACCGAGGAGCTGCGGGTGCCTCGCCTCCCACCTGCGCAAGCTGCGTAAGCGG
CTCCTCCGCGATGCCGATGACCTGCAGAAGCGCCTGGCAGTGTACCAGGCCGGGGCCCGC
GAGGGCGCCGAGCGCGGCTCAGCGCCATCCGCGAGCGCCTGGGGCCCTGGTGGAAACAG
GGCCGCGTGGCGGCCGCCACTGTGGGCTCCCTGGCCGGCCAGCCGCTACAGGAGCGGGCC
CAGGCCTGGGGCGAGCGGCTGCGCGCGCGGATGGAGGAGATGGGCAGCCGGACCCGCGAC
CGCCTGGACGAGGTGAAGGAGCAGGTGGCGGAGGTGCGCGCCAAGCTGGAGGAGCAGGCC
CAGCAGATACGCCTGCAGGCCGAGGCCTTCCAGGCCCGCCTCAAGAGCTGGTTCGAGCCC
CTGGTGGAAAGACATGCAGCGCCAGTGGGCCGGCTGGTGGAGAAGGTGCAGGCTGCCGTG
GGCACCAGCGCCGCCCTGTGCCAGCGACAATCACTGAACGCCGAAGCCTGCAGCCATG
CGACCCACGCCACCCCGTGCCTCCTCGCTCCGCGCAGCCTGCAGCGGGAGACCCTGTCC
CCGCCCCAGCCGTCTCCTGGGGTGGACCCTAGTTTAATAAAGATTACCAAGTTTCACG
CAAAC

Restriction Sites: Please inquire
ACCN: NM_000041



[View online »](#)

OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	NM_000041.2 , NP_000032.1
RefSeq Size:	1223 bp
RefSeq ORF:	954 bp
Locus ID:	348
UniProt ID:	P02649
Cytogenetics:	19q13.32
Domains:	Apolipoprotein
Protein Families:	Adult stem cells, Druggable Genome, Secreted Protein, Stem cell - Pluripotency
Protein Pathways:	Alzheimer's disease
Gene Summary:	<p>The protein encoded by this gene is a major apoprotein of the chylomicron. It binds to a specific liver and peripheral cell receptor, and is essential for the normal catabolism of triglyceride-rich lipoprotein constituents. This gene maps to chromosome 19 in a cluster with the related apolipoprotein C1 and C2 genes. Mutations in this gene result in familial dysbetalipoproteinemia, or type III hyperlipoproteinemia (HLP III), in which increased plasma cholesterol and triglycerides are the consequence of impaired clearance of chylomicron and VLDL remnants. [provided by RefSeq, Jun 2016]</p> <p>Transcript Variant: This variant (2) contains an alternate 5' terminal exon, and it thus differs in the 5' UTR and initiates translation at a downstream in-frame start codon, compared to variant 1. The encoded isoform (b) is shorter at the N-terminus, compared to isoform a. Variants 2, 3, 4 and 5 all encode isoform b.</p>