

## Product datasheet for **SC319630**

### GCAT (NM\_014291) Human Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	GCAT (NM_014291) Human Untagged Clone
Tag:	Tag Free
Symbol:	GCAT
Synonyms:	KBL
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC (PS100020)
E. coli Selection:	Ampicillin (100 ug/mL)

**Fully Sequenced ORF:** >OriGene sequence for NM\_014291.2  
GCTCGGGCGAGGTAGGAGCGATGTGGCCTGGGAACGCCTGGCGCGCCGCACTCTTCTGGG  
TGCCCCGCGGCCCGCCGCACAGTCAGCGCTGGCCAGCTGCGTGGCATTCTGGAGGGG  
AGCTGGAAGGCATCTGCGGAGCTGGCACTTGAAGAGTGAGCGGTGCATCACGTCCCGTC  
AGGGGCCGCACATCCGCGTGGACGGCGTCTCCGAGGAATCCTTAACTTCTGTGCCAACA  
ACTACCTGGGCCTGAGCAGCCACCCTGAGGTGATCCAGGCAGGTCTGCAGGCTCTGGAGG  
AGTTTGGAGCTGGCCTCAGCTCTGTCCGCTTTATCTGTGGAACCCAGAGCATCCACAAGA  
ATCTAGAAGCAAAAATAGCCCGTTCCACCAGCGGGAGGATGCCATCCTCTATCCCAGCT  
GTTATGACGCCAACGCCGCTCTTTGAGGCCCTGCTGACCCAGAGGACGCAGTCTGT  
CGGACGAGCTGAACCATGCCTCCATCATCGACGGCATCCGGCTGTGCAAGGCCACAAGT  
ACCGCTATCGCCACCTGGACATGGCCGACCTAGAAGCCAAGCTGCAGGAGGCCAGAAGC  
ATCGGCTGCGCCTGGTGGCCACTGATGGGCCTTTTCCATGGATGGCGACATCGCACCCC  
TGCAGGAGATCTGCTGCCTCGCCTCTAGATATGGTGCCTGGTCTTCATGGATGAATGCC  
ATGCCACTGGCTTCTGGGGCCACAGGACGGGGCACAGATGAGCTGCTGGGTGTGATGG  
ACCAGGTCAACATCACTCACTCCACCCTGGGGAAGGCCCTGGGTGGAGCATCAGGGGGCT  
ACACGACAGGGCCTGGGCCCTGGTGTCCCTGCTGCGGCAGCGCGCCCGCCATACCTCT  
TCTCCAACAGTCTGCCACCTGCTGTCGTTGGCTGCGCCTCCAAGGCCCTAGATCTGCTGA  
TGGGAGTAACACCATTTGCCAGTCTATGGCTGCCAAGACCCAGAGGTTCCGTAGTAAGA  
TGAAGCTGCTGGCTTCACTATCTCGGGAGCCAGTCACCCCATCTGCCCTGTGATGCTGG  
GTGATGCCCGGCTGGCCTCTCGCATGGCGGATGACATGCTGAAGAGAGGCATCTTTGTCA  
TCGGGTTCACTACCCCGTGGTCCCAAGGGCAAGGCCTGGATCCGGGTACAGATCTCAG  
CAGTGCATAGCGAGGAAGACATTGACCGCTGCGTGGAGGCCTTCGTGGAAGTGGGGCGAC  
TGCACGGGGCACTGCCCTGAGCTCTGGTAAGGACGAGAAGAGCCAAGGTCGCTGCTG  
CCACAGGGTCAAAGGAGGTTTTCGATCAGCCCAGACCAGAGGCTCTGAGCCCTGAACCAA  
AGTCCCAGAGCTGGGCTGGGACGTGACCTGTGCTGAGGGCTGTGAGAATGTGAAACAACA  
GTGTGAAAATTGGCTGTGCCAAAAAAAAAAAAAAAAAAAAAAAAAAAAA



[View online »](#)

<b>Restriction Sites:</b>	Please inquire
<b>ACCN:</b>	NM_014291
<b>OTI Disclaimer:</b>	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
<b>OTI Annotation:</b>	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"> <li>1. Centrifuge at 5,000xg for 5min.</li> <li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>3. Close the tube and incubate for 10 minutes at room temperature.</li> <li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
<b>RefSeq:</b>	<u><a href="#">NM_014291.2</a></u> , <u><a href="#">NP_055106.1</a></u>
<b>RefSeq Size:</b>	1470 bp
<b>RefSeq ORF:</b>	1260 bp
<b>Locus ID:</b>	23464
<b>UniProt ID:</b>	<u><a href="#">O75600</a></u>
<b>Cytogenetics:</b>	22q13.1
<b>Domains:</b>	aminotran_1_2
<b>Protein Pathways:</b>	Glycine, serine and threonine metabolism

**Gene Summary:**

The degradation of L-threonine to glycine consists of a two-step biochemical pathway involving the enzymes L-threonine dehydrogenase and 2-amino-3-ketobutyrate coenzyme A ligase. L-Threonine is first converted into 2-amino-3-ketobutyrate by L-threonine dehydrogenase. This gene encodes the second enzyme in this pathway, which then catalyzes the reaction between 2-amino-3-ketobutyrate and coenzyme A to form glycine and acetyl-CoA. The encoded enzyme is considered a class II pyridoxal-phosphate-dependent aminotransferase. Alternate splicing results in multiple transcript variants. A pseudogene of this gene is found on chromosome 14. [provided by RefSeq, Jan 2010]

Transcript Variant: This variant (2) differs in the 3' UTR and uses an alternate splice site in the coding region, compared to variant 1. The encoded isoform (2) is shorter than isoform 1.

Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.