

## Product datasheet for **SC319936**

### SERF2 (NM\_001018108) Human Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	SERF2 (NM_001018108) Human Untagged Clone
Tag:	Tag Free
Symbol:	SERF2
Synonyms:	4F5REL; FAM2C; H4F5REL; HsT17089
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC (PS100020)
E. coli Selection:	Ampicillin (100 ug/mL)
Fully Sequenced ORF:	>NCBI ORF sequence for NM_001018108, the custom clone sequence may differ by one or more nucleotides

ATGACCCGCGGTAACCAGCGTGAGCTCGCCCGCCAGAAGAATATGAAAAAGCAGAGCGACTCGGTTAAGG  
 GAAAGCGCCGAGATGACGGGCTTTCTGCTGCCGCCGCAAGCAGAGGGACTCGGAGATCATGCAGCAGAA  
 GCAGAAAAAGGCAAACGAGAAGAAGGAGGAACCCAAGTAG

Restriction Sites:	Please inquire
ACCN:	NM_001018108
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).


[View online »](#)

<b>Reconstitution Method:</b>	<ol style="list-style-type: none"> <li>1. Centrifuge at 5,000xg for 5min.</li> <li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>3. Close the tube and incubate for 10 minutes at room temperature.</li> <li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
<b>RefSeq:</b>	<u>NM_001018108.1, NP_001018118.1</u>
<b>RefSeq Size:</b>	530 bp
<b>RefSeq ORF:</b>	180 bp
<b>Locus ID:</b>	10169
<b>UniProt ID:</b>	<u>P84101</u>
<b>Cytogenetics:</b>	15q15.3
<b>Gene Summary:</b>	<p>Positive regulator of amyloid protein aggregation and proteotoxicity (PubMed:20723760). Induces conformational changes in amyloid proteins, such as HTT, driving them into compact formations preceding the formation of aggregates (PubMed:20723760).[UniProtKB/Swiss-Prot Function]</p> <p>Transcript Variant: This variant (3) differs in the 5' UTR, and uses an alternate splice site that causes a frameshift in the 3' coding region, compared to variant 1. The encoded isoform (c) has a distinct and shorter C-terminus, compared to isoform a. Both variants 3 and 4 encode isoform c. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.</p>