

## **Product datasheet for SC320112**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Mimitin (NDUFAF2) (NM\_174889) Human Untagged Clone

#### **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** Mimitin (NDUFAF2) (NM\_174889) Human Untagged Clone

Tag: Tag Free
Symbol: Mimitin

Synonyms: B17.2L; MC1DN10; mimitin; MMTN; NDUFA12L

Mammalian Cell

Selection:

None

Vector: pCMV6-XL5

E. coli Selection: Ampicillin (100 ug/mL)

Fully Sequenced ORF: >OriGene sequence for NM\_174889.2

**Restriction Sites:** Please inquire ACCN: NM\_174889

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.





### Mimitin (NDUFAF2) (NM\_174889) Human Untagged Clone - SC320112

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 174889.2</u>, <u>NP 777549.1</u>

 RefSeq Size:
 650 bp

 RefSeq ORF:
 510 bp

 Locus ID:
 91942

 UniProt ID:
 Q8N183

 Cytogenetics:
 5q12.1

Gene Summary: NADH:ubiquinone oxidoreductase (complex I) catalyzes the transfer of electrons from NADH

to ubiquinone (coenzyme Q) in the first step of the mitochondrial respiratory chain, resulting in the translocation of protons across the inner mitochondrial membrane. This gene encodes

a complex I assembly factor. Mutations in this gene cause progressive encephalopathy

resulting from mitochondrial complex I deficiency. [provided by RefSeq, Jul 2008]