

Product datasheet for **SC320823**

HBA1 (NM_000558) Human Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: HBA1 (NM_000558) Human Untagged Clone
Tag: Tag Free
Symbol: HBA1
Synonyms: ECYT7; HBA-T3; HBH; METHBA
Mammalian Cell Selection: Neomycin
Vector: pCMV6-AC (PS100020)
E. coli Selection: Ampicillin (100 ug/mL)

Fully Sequenced ORF: >OriGene sequence for NM_000558.3
ACTTTCTGGTCCCCACAGACTCAGAGAGAACCCACCATGGTGCTGTCTCCTGCCGACAA
GACCAACGTCAAGGCCGCTGGGGTAAGGTCGGCGCGCACGCTGGCGAGTATGGTGCGGA
GGCCCTGGAGAGGATGTTCTGTCTTCCCCACCACCAAGACCTACTTCCGCACTTCGA
CCTGAGCCACGGCTCTGCCAGGTTAAGGGCCACGGCAAGAAGGTGGCCGACGCGTGAC
CAACGCGTGGCGCACGTGGACGACATGCCAACGCGCTGTCCGCCCTGAGCGACCTGCA
CGCGCACAAAGCTTCGGGTGGACCCGGTCAACTTCAAGCTCCTAAGCCACTGCCTGCTGGT
GACCCTGGCCGCCACCTCCCCCGGAGTTCACCCCTGCGGTGCACGCTCCCTGGACAA
GTTCTGGCTTCTGTGAGCACCGTGCTGACCTCAAATACCGTTAAGCTGGAGCCTCGGT
GGCCATGCTTCTGCCCCTTGGGCTCCCCCAGCCCTCCTCCCCTTCTGCACCCGTA
CCCCCGTGGTCTTGAATAAAGTCTGAGTGGGCGCAAAAAAAAAAAAAAAAAAAAAA
AACAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

Restriction Sites: Please inquire

ACCN: NM_000558

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.



[View online »](#)

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_000558.3, NP_000549.1</u>
RefSeq Size:	576 bp
RefSeq ORF:	429 bp
Locus ID:	3039
UniProt ID:	<u>P69905</u>
Cytogenetics:	16p13.3
Domains:	globin
Gene Summary:	<p>The human alpha globin gene cluster located on chromosome 16 spans about 30 kb and includes seven loci: 5'- zeta - pseudozeta - mu - pseudoalpha-1 - alpha-2 - alpha-1 - theta - 3'. The alpha-2 (HBA2) and alpha-1 (HBA1) coding sequences are identical. These genes differ slightly over the 5' untranslated regions and the introns, but they differ significantly over the 3' untranslated regions. Two alpha chains plus two beta chains constitute HbA, which in normal adult life comprises about 97% of the total hemoglobin; alpha chains combine with delta chains to constitute HbA-2, which with HbF (fetal hemoglobin) makes up the remaining 3% of adult hemoglobin. Alpha thalassemias result from deletions of each of the alpha genes as well as deletions of both HBA2 and HBA1; some nondeletion alpha thalassemias have also been reported. [provided by RefSeq, Jul 2008]</p>