

Product datasheet for SC320876

PSMD7 (NM 002811) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: PSMD7 (NM_002811) Human Untagged Clone

Tag: Tag Free
Symbol: PSMD7

Synonyms: MOV34; P40; Rpn8; S12

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-AC (PS100020)E. coli Selection:Ampicillin (100 ug/mL)

Fully Sequenced ORF: >OriGene sequence for NM_002811.3

GGGGCGAGGGCTGACGAACCGGAAGAAGAGGAACTGGGCCTGAAAGGGTACCGGTGACCG CTACTGCTGCCGGTGTTTGCGTGTGGCAGGGAGCCAGGCCTGGCGAGCGGGGTGTGTCGC GATGCCGGAGCTGGCAGTGCAGAAGGTGGTGGTCCACCCCCTGGTGCTCAGTGTGGT GGATCATTTCAACCGAATCGGCAAGGTTGGAAACCAGAAGCGTGTTGTTGGTGTGCTTTT GGGGTCATGGCAAAAGAAAGTACTTGATGTATCGAACAGTTTTGCAGTTCCTTTTGATGA AGATGACAAAGACGATTCTGTATGGTTTTTAGACCATGATTATTTGGAAAACATGTATGG AATGTTTAAGAAAGTCAATGCCAGGGAAAGAATAGTTGGCTGGTACCACACAGGCCCTAA ACTACACAAGAATGACATTGCCATCAACGAACTCATGAAAAGATACTGTCCTAATTCCGT ATTGGTCATCATTGATGTGAAGCCGAAGGACCTAGGGCTGCCTACAGAAGCGTACATTTC AGTGGAAGAAGTCCATGATGATGGAACTCCAACCTCGAAAACATTTGAACACGTGACCAG TGAAATTGGAGCAGAGGAAGCTGAGGAAGTTGGAGTTGAACACTTGTTACGAGATATCAA AGACACGACGGTGGGCACTCTGTCCCAGCGGATCACAAACCAGGTCCATGGTTTGAAGGG ACTGAACTCCAAGCTTCTGGATATCAGGAGCTACCTGGAAAAAGTCGCCACAGGCAAGCT GCCCATCAACCACCAGATCATCTACCAGCTGCAGGACGTCTTCAACCTGCTGCCAGATGT CAGCCTGCAGGAGTTCGTCAAGGCCTTTTACCTGAAGACCAATGACCAGATGGTGGTAGT GTACTTGGCCTCGCTGATCCGTTCCGTGGTCGCCCTGCACAACCTCATCAACAACAAGAT GAAAGAGGACAAGGAGAAAGATAAAGATAAGGAAAAGAGTGATGTAAAGAAAGAGAGAGAA AAAGGAGAAAAAGTAAAACATGTATTAAATAGCTTTTTTAATTTGTAAATTAAAATCTTA

CAAACCAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

Restriction Sites: Please inquire ACCN: NM_002811



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

PSMD7 (NM_002811) Human Untagged Clone - SC320876

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

> point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the

expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

NM 002811.3, NP 002802.2 RefSeq:

RefSeg Size: 1678 bp RefSeq ORF: 975 bp Locus ID: 5713

Cytogenetics: 16q23.1 **Domains:** JAB MPN

UniProt ID:

Protein Families: Druggable Genome, Protease

P51665

Protein Pathways: Proteasome

Gene Summary: The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure

> composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a non-ATPase subunit of the 19S regulator. A pseudogene has been identified on chromosome 17. [provided by RefSeq, Jul 2008]