

Product datasheet for SC320947

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

H4C11 (NM_021968) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: H4C11 (NM_021968) Human Untagged Clone

Tag: Tag Free Symbol: H4C11

Synonyms: dJ160A22.2; H4-16; H4/e; H4C1; H4C2; H4C3; H4C4; H4C5; H4C6; H4C8; H4C9; H4C12; H4C13;

H4C14; H4C15; H4F2iv; H4FE; HIST1H4I

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-AC (PS100020)E. coli Selection:Ampicillin (100 ug/mL)

Fully Sequenced ORF: >OriGene sequence for NM_021968.3

AAAAA

Restriction Sites: Please inquire **ACCN:** NM 021968

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 021968.3</u>, <u>NP 068803.1</u>

 RefSeq Size:
 356 bp

 RefSeq ORF:
 312 bp

 Locus ID:
 8363

 UniProt ID:
 P62805

 Cytogenetics:
 6p22.1

Protein Pathways: Systemic lupus erythematosus

Gene Summary: Histones are basic nuclear proteins that are responsible for the nucleosome structure of the

chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H4 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the small histone gene

cluster on chromosome 6p22-p21.3. [provided by RefSeq, Aug 2015]