

Product datasheet for SC321044

Caspase 1 (CASP1) (NM 033293) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Caspase 1 (CASP1) (NM_033293) Human Untagged Clone

Tag: Tag Free

Symbol: Caspase 1

Synonyms: ICE; IL1BC; P45

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-AC (PS100020)E. coli Selection:Ampicillin (100 ug/mL)

Fully Sequenced ORF: >OriGene sequence for NM_033293.2

GAGAAAAGCCATGGCCGACAAGGTCCTGAAGGAGAAGAGAAAGCTGTTTATCCGTTCCAT GGGTGAAGCTCCTCAGGCAGTGCAGGACAACCCAGCTATGCCCACATCCTCAGGCTCAGA AGGGAATGTCAAGCTTTGCTCCCTAGAAGAAGCTCAAAGGATATGGAAACAAAAGTCGGC AGAGATTTATCCAATAATGGACAAGTCAAGCCGCACACGTCTTGCTCTCATTATCTGCAA TGAAGAATTTGACAGTATTCCTAGAAGAACTGGAGCTGAGGTTGACATCACAGGCATGAC CATGACTACAGAGCTGGAGGCATTTGCACACCGCCCAGAGCACAAGACCTCTGACAGCAC GTTCCTGGTGTTCATGTCTCATGGTATTCGGGAAGGCATTTGTGGGAAGAACACTCTGA GCAAGTCCCAGATATACTACAACTCAATGCAATCTTTAACATGTTGAATACCAAGAACTG CCCAAGTTTGAAGGACAAACCGAAGGTGATCATCATCCAGGCCTGCCGTGGTGACAGCCC TGGTGTGGTGTGGTTTAAAGATTCAGTAGGAGTTTCTGGAAACCTATCTTTACCAACTAC AGAAGAGTTTGAGGATGATGCTATTAAGAAAGCCCACATAGAGAAGGATTTTATCGCTTT CTGCTCTTCCACACCAGATAATGTTTCTTGGAGACATCCCACAATGGGCTCTGTTTTTAT TGGAAGACTCATTGAACATATGCAAGAATATGCCTGTTCCTGTGATGTGGAGGAAATTTT CCGCAAGGTTCGATTTTCATTTGAGCAGCCAGATGGTAGAGCGCAGATGCCCACCACTGA AAGAGTGACTTTGACAAGATGTTTCTACCTCTTCCCAGGACATTAAAATAAGGAAACTGT ATGAATGTCTGTGGGCAGGAAGTGAAGAGATCCTTCTGTAAAGGTTTTTGGAATTATGTC

AAAAAAAAA

Restriction Sites: Please inquire **ACCN:** NM 033293



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Caspase 1 (CASP1) (NM_033293) Human Untagged Clone - SC321044

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 033293.2</u>, <u>NP 150635.1</u>

RefSeq Size:1085 bpRefSeq ORF:936 bp

UniProt ID: P29466

Locus ID:

Cytogenetics: 11q22.3

Protein Families: Druggable Genome, Protease

834

Protein Pathways: Amyotrophic lateral sclerosis (ALS), Cytosolic DNA-sensing pathway, NOD-like receptor

signaling pathway



Gene Summary:

This gene encodes a protein which is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce 2 subunits, large and small, that dimerize to form the active enzyme. This gene was identified by its ability to proteolytically cleave and activate the inactive precursor of interleukin-1, a cytokine involved in the processes such as inflammation, septic shock, and wound healing. This gene has been shown to induce cell apoptosis and may function in various developmental stages. Studies of a similar gene in mouse suggest a role in the pathogenesis of Huntington disease. Alternative splicing results in transcript variants encoding distinct isoforms. [provided by RefSeq, Mar 2012]

Transcript Variant: This variant (gamma) has multiple differences in the coding region but the translation remains in frame, compared to variant alpha. Variant gamma encodes a protein lacking an internal segment, as compared to isoform alpha. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.