

# **Product datasheet for SC321047**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## CCL4L1 (NM\_207007) Human Untagged Clone

### **Product data:**

**Product Type:** Expression Plasmids

Product Name: CCL4L1 (NM\_207007) Human Untagged Clone

Tag: Tag Free Symbol: CCL4L1

Synonyms: AT744.2; CCL4L; LAG-1; LAG1; MIP-1-beta; SCYA4L; SCYA4L1; SCYA4L2

**Mammalian Cell** 

Selection:

Neomycin

Vector:pCMV6-AC (PS100020)E. coli Selection:Ampicillin (100 ug/mL)

Fully Sequenced ORF: >OriGene sequence for NM\_207007.2

AAAAAAAAAAAAAAAAAAAAA

**Restriction Sites:** Please inquire **ACCN:** NM 207007

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.



### CCL4L1 (NM\_207007) Human Untagged Clone - SC321047

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** <u>NM 207007.2</u>, <u>NP 996890.1</u>

 RefSeq Size:
 685 bp

 RefSeq ORF:
 279 bp

 Locus ID:
 388372

 UniProt ID:
 P13236

 Cytogenetics:
 17q12

**Protein Families:** Druggable Genome, Transmembrane

**Protein Pathways:** Chemokine signaling pathway, Cytokine-cytokine receptor interaction, Cytosolic DNA-sensing

pathway

**Gene Summary:** This gene is one of several cytokine genes that are clustered on the q-arm of chromosome 17.

Cytokines are a family of secreted proteins that function in inflammatory and

immunoregulatory processes. The protein encoded by this family member is similar to the chemokine (C-C motif) ligand 4 product, which inhibits HIV entry by binding to the cellular receptor CCR5. The copy number of this gene varies among individuals, where most individuals have one to five copies. Alternative splicing of this gene results in multiple

transcript variants. [provided by RefSeq, Apr 2014]

Transcript Variant: This variant (CCL4L) represents the longer transcript and encodes the

supported protein.