

Product datasheet for **SC322038**

HSPC142 (BABAM1) (NM_001033549) Human Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: HSPC142 (BABAM1) (NM_001033549) Human Untagged Clone
Tag: Tag Free
Symbol: HSPC142
Synonyms: C19orf62; HSPC142; MERIT40; NBA1
Mammalian Cell Selection: Neomycin
Vector: pCMV6-AC (PS100020)
E. coli Selection: Ampicillin (100 ug/mL)

Fully Sequenced ORF: >OriGene sequence for SC322038
CGGCTTCTAGTGTGAGTCGGCGGCTGATTTAGAAGGAGTTTCAGGCTACGGTGAGCCGAAG
GGGAGGATTCCTGGCTTCCCTGTCCGTGTTCCATCTAGCCACACAGGAGCCATGGAAGTG
GCAGAGCCCAGCAGCCCCACTGAAGAGGAGGAGGAGGAAGAGGAGCACTCGGCAGAGCCT
CGGCCCCGCACTCGCTCCAATCCTGAAGGGCTGAGGACCGGGCAGTAGGGGCACAGGCC
AGCGTGGGACGCCAGCGAGGGTGAAGGTGAGGCCGCCAGTGCTGATGATGGGAGCCTC
AACACTTCAGGAGCCGGCCCTAAGTCTGGCAGGTGCCCCGCCAGCCCCTGAGGTCCAA
ATTCGGACACCAAGGGTCAACTGTCCAGAGAAAGTGATTATCTGCCTGGACCTGTCAGAG
GAAATGCACTGCCAAAGCTGGAGTCGTTCAACGGCTCCAAAACCAACGCCCTCAATGTC
TCCCAGAAGATGATTGAGATGTTTCGTGCGGACAAAACACAAGATCGACAAAAGCCACGAG
TTTGCCTGGTGGTGGTGAACGATGACACGGCCTGGCTGTCTGGCCTGACCTCCGACCCC
CGCGAGCTCTGTAGCTGCCTCTATGATCTGGAGACGGCCTCCTGTTCCACCTTCAATCTG
GAAGGACTTTTCAGCCTCATCCAGCAGAAAAGTGAAGTTCGGTCCAGAGAACGTGCAG
ACGATTCACCCCGCATATGTGGTCCGCACCATCCTTGTCTACAGCCGTCCACCTTGCCAG
CCCCAGTTCTCCTTGACGGAGCCCATGAAGAAAATGTTCCAGTGCCCATATTTCTTCTTT
GACGTTGTTTACATCCACAATGGCACTGAGGAGAAGGAGGAGGAGATGAGTTGGAAGGAT
ATGTTTGCCTTCATGGGCAGCCTGGATACCAAGGGTACCAGCTACAAGTATGAGGTGGCA
CTGGCTGGGCCAGCCCTGGAGTTGCACAACCTGCATGGCGAAACTGTTGGCCACCCCTG
CAGCGCCCTTGGCAGAGCCATGCTTCTACAGCCTGCTGGAGGAGGAGGATGAAGCCATT
GAGGTTGAGGCCACTGTCTGAACCATCCCTGTACATCTGCACCTTCTTGTGCAAGGAAGT
CCTTGGCCTAAAGCCTTGGTTCTCAAAGTGGGTTCCCTGGGACCTCCGGGTGGGGGGT
TCCAGGAGGCACGTAGGGTACCTTGCAGGGTCTAGGAGGAAACCCAGGATTCCAGGAG
GGATCCCAGGAAGTGTGGGCACCCATTTTCTGTGTCTCCAGCCATTTCCACTCCTAGT
TTGTCATGGATAATTTTGTCTCCCTGTGTGATTTTGGCCATCAAATAAAAAATTTGA
GACTCGTTAAAAAAAAAAAAAAAAAAAA

Restriction Sites: Please inquire



[View online >](#)

ACCN:	NM_001033549
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001033549.1, NP_001028721.1</u>
RefSeq Size:	1451 bp
RefSeq ORF:	990 bp
Locus ID:	29086
UniProt ID:	<u>Q9NWX8</u>
Cytogenetics:	19p13.11

Gene Summary:

Component of the BRCA1-A complex, a complex that specifically recognizes 'Lys-63'-linked ubiquitinated histones H2A and H2AX at DNA lesions sites, leading to target the BRCA1-BARD1 heterodimer to sites of DNA damage at double-strand breaks (DSBs). The BRCA1-A complex also possesses deubiquitinase activity that specifically removes 'Lys-63'-linked ubiquitin on histones H2A and H2AX. In the BRCA1-A complex, it is required for the complex integrity and its localization at DSBs. Component of the BRISC complex, a multiprotein complex that specifically cleaves 'Lys-63'-linked ubiquitin in various substrates (PubMed:24075985, PubMed:26195665). In these 2 complexes, it is probably required to maintain the stability of BABAM2 and help the 'Lys-63'-linked deubiquitinase activity mediated by BRCC3/BRCC36 component. The BRISC complex is required for normal mitotic spindle assembly and microtubule attachment to kinetochores via its role in deubiquitinating NUMA1 (PubMed:26195665). Plays a role in interferon signaling via its role in the deubiquitination of the interferon receptor IFNAR1; deubiquitination increases IFNAR1 activity by enhancing its stability and cell surface expression (PubMed:24075985). Down-regulates the response to bacterial lipopolysaccharide (LPS) via its role in IFNAR1 deubiquitination (PubMed:24075985).[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (1) is the longest transcript and encodes the longer isoform (1). Variants 1, 2 and 3 encode the same isoform (1).