

Product datasheet for SC323062

CYP2C18 (NM_001128925) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	CYP2C18 (NM_001128925) Human Untagged Clone
Tag:	Tag Free
Symbol:	CYP2C18
Synonyms:	CPCI; CYP2C; CYP2C17; P450-6B/29C; P450IIC17
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC323062 representing NM_001128925. Blue=Insert sequence Red=Cloning site Green=Tag(s)


```
GCTCGTTT TAGTGAACCGTCAGAATTTTGT AATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTG
GATCCGGTACCGAGGAGATCTGCCGCC CGATCGCC
ATGGATCCAGCTGTGGCTCTGGTCTGTCTCTCCTGTTTGTTCCTTTCACTCTGGAGGCAGAGC
TCTGGAAGAGGGAGGCTCCCGTCTGGCCCCACTCCTCTCCCGATTATTGGAAATATCCTGCAGTTAGAT
GTTAAGGACATGAGCAAATCCTTAACCAATTTCTCAAAGTCTATGGCCCTGTGTTCACTGTGATTTTT
GGCCTGAAGCCCATTGTGGTGTGCATGGATATGAAGCAGTGAAGGAGGCCCTGATTGATCATGGAGAG
GAGTTTTCTGGAAGAGGAAGTTTTCCAGTGGCTGAAAAAGTTAACAAAGGACTTGAATCCTTTTCAGC
AATGAAAAGAGATGGAAGGAGATCCGGCGTTTCTGCCTCATGACTCTGCGGAATTTGGGATGGGAAG
AGGAGCATCGAGGACCGTGTCAAGAGGAAGCCCGCTGCCTTGTGGAGGAGTTGAGAAAAACCAATGCC
TCAACCCTGTGATCCCACTTTTCATCCTGGGCTGTGCTCCCTGCAATGTGATCTGCTCTGTTATTTCCAT
GATCGATTTGATTATAAAGATCAGAGGTTTCTTAACCTGATGGAAAAATCAATGAAAACCTCAGGATT
CTGAGCTCTCCATGGATCCAGGAAAAGCACAATCAACAGTCTGAATTTACTGTTGAAAGCTTGATAGCC
ACTGTAAGTATGATGTTTGGGCTGGAACAGAGACAACGAGCACCCTCTGAGATATGGACTCCTGCTC
CTGCTGAAGTACCCAGAGGTCACAGCTAAAGTCCAGGAAGAGATTGAATGTGTAGTTGGCAGAAACCGG
AGCCCCGTATGCAGGACAGGAGTCACATGCCCTACACAGATGCTGTGGTGCACGAGATCCAGAGATAC
ATTGACCTCCTCCCACCAACCTGCCCATGCAGTACCTGTGATGTTAAATTCAAAACCTACCTCATC
CCCAAGGGCAGCACCATAATAACATCCCTGACTTCTGTGCTGCACAATGACAAAGAATCCCAACCCA
GAGATGTTTGACCCTGGCACTTTCTGGATAAAGAGTGGCAACTTTAAGAAAAGTACTACTTTCATGCCT
TTCTCAGCAGGAAAACGGATGTGTATGGGAGAGGGCTGGCCCGCATGGAGCTGTTTTATTCTGACC
ACCATTTTGCAGAACTTTAACCTGAAATCTCAGGTTGACCCAAAGGATATTGACATCACCCCATTTGCC
AATGCATTTGGTCGTGTGCCACCCTGTACCAGCTCTGCTTCATTCTGTCTGA
ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
```



[View online »](#)

Restriction Sites: SgfI-MluI

Plasmid Map:



ACCN: NM_001128925

Insert Size: 1296 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001128925.1](#)

RefSeq Size: 2385 bp

RefSeq ORF: 1296 bp

Locus ID: 1562

UniProt ID:	<u>P33260</u>
Cytogenetics:	10q23.33
Protein Families:	Druggable Genome, P450
Protein Pathways:	Arachidonic acid metabolism, Drug metabolism - cytochrome P450, Linoleic acid metabolism, Metabolic pathways, Metabolism of xenobiotics by cytochrome P450, Retinol metabolism
MW:	48.8 kDa
Gene Summary:	<p>This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum but its specific substrate has not yet been determined. The gene is located within a cluster of cytochrome P450 genes on chromosome 10q24. An additional gene, CYP2C17, was once thought to exist; however, CYP2C17 is now considered an artefact based on a chimera of CYP2C18 and CYP2C19. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]</p> <p>Transcript Variant: This variant (2) lacks a coding exon in the middle region, but the reading frame is not affected, as compared to variant 1. The resulting isoform (2) lacks an internal segment, as compared to isoform 1.</p>