

## **Product datasheet for SC323348**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## S6K1 (RPS6KB1) (NM\_003161) Human Untagged Clone

## **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** S6K1 (RPS6KB1) (NM\_003161) Human Untagged Clone

Tag: Tag Free Symbol: S6K1

**Synonyms:** p70 S6KA; p70(S6K)-alpha; p70-alpha; p70-S6K; PS6K; S6K-beta-1; S6K1; STK14A

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)



Fully Sequenced ORF: >SC323348 representing NM\_003161.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

GCTCGTTTAGTGAACCGTCAGAATTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

ATGAGGCGACGAAGGAGGCGGACGGCTTTTACCCAGCCCCGGACTTCCGAGACAGGGAAGCTGAGGAC ATGGCAGGAGTGTTTGACATAGACCTGGACCAGCCAGAGGACGCGGGCTCTGAGGATGAGCTGGAGGAG GGGGGTCAGTTAAATGAAAGCATGGACCATGGGGGAGTTGGACCATATGAACTTGGCATGGAACATTGT GAGAAATTTGAAATCTCAGAAACTAGTGTGAACAGAGGGCCAGAAAAAATCAGACCAGAATGTTTTGAG CTACTTCGGGTACTTGGTAAAGGGGGCTATGGAAAGGTTTTTCAAGTACGAAAAGTAACAGGAGCAAAT ACTGGGAAAATATTTGCCATGAAGGTGCTTAAAAAGGCAATGATAGTAAGAAATGCTAAAGATACAGCT CATACAAAAGCAGAACGGAATATTCTGGAGGAAGTAAAGCATCCCTTCATCGTGGATTTAATTTATGCC TTTCAGACTGGTGGAAAACTCTACCTCATCCTTGAGTATCTCAGTGGAGAGAACTATTTATGCAGTTA GAAAGAGAGGGAATATTTATGGAAGACACTGCCTGCTTTTACTTGGCAGAAATCTCCATGGCTTTGGGG CATTTACATCAAAAGGGGATCATCTACAGAGACCTGAAGCCGGAGAATATCATGCTTAATCACCAAGGT TGTGGAACAATAGAATACATGGCCCCTGAAATCTTGATGAGAAGTGGCCACAATCGTGCTGTGGATTGG TGGAGTTTGGGAGCATTAATGTATGACATGCTGACTGGAGCACCCCCATTCACTGGGGAGAATAGAAAG AAAACAATTGACAAAATCCTCAAATGTAAACTCAATTTGCCTCCCTACCTCACACAAGAAGCCAGAGAT CTGCTTAAAAAGCTGCTGAAAAGAAATGCTGCTTCTCGTCTGGGAGCTGGTCCTGGGGACGCTGGAGAA GTTCAAGCTCATCCATTCTTTAGACACATTAACTGGGAAGAACTTCTGGCTCGAAAGGTGGAGCCCCCC TTTAAACCTCTGTTGCAATCTGAAGAGGATGTAAGTCAGTTTGATTCCAAGTTTACACGTCAGACACCT GTCGACAGCCCAGATGACTCAACTCTCAGTGAAAGTGCCAATCAGGTCTTTCTGGGTTTTACATATGTG GCTCCATCTGTACTTGAAAGTGTGAAAGAAAAGTTTTCCTTTGAACCAAAAATCCGATCACCTCGAAGA TTTATTGGCAGCCCACGAACACCTGTCAGCCCAGTCAAATTTTCTCCTGGGGATTTCTGGGGAAGAGGT GCTTCGGCCAGCACAGCAAATCCTCAGACACCTGTGGAATACCCAATGGAAACAAGTGGCATAGAGCAG ATGGATGTGACAATGAGTGGGGAAGCATCGGCACCACTTCCAATACGACAGCCGAACTCTGGGCCATAC AAAAAACAAGCTTTTCCCATGATCTCCAAACGGCCAGAGCACCTGCGTATGAATCTA<mark>TGA</mark>

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGGATCTGGCAGCAAATGATATCCTGGAT

TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

Restriction Sites: Sgfl-Mlul
ACCN: NM\_003161
Insert Size: 1578 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts

of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by

calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>

ORÏGENE

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** <u>NM 003161.3</u>

 RefSeq Size:
 5368 bp

 RefSeq ORF:
 1578 bp

 Locus ID:
 6198

 UniProt ID:
 P23443

 Cytogenetics:
 17q23.1

**Domains:** pkinase, S\_TK\_X, TyrKc, S\_TKc

**Protein Families:** Druggable Genome, Protein Kinase

Protein Pathways: Acute myeloid leukemia, ErbB signaling pathway, Fc gamma R-mediated phagocytosis, Insulin

signaling pathway, mTOR signaling pathway, TGF-beta signaling pathway

**MW:** 59.1 kDa

**Gene Summary:** This gene encodes a member of the ribosomal S6 kinase family of serine/threonine kinases.

The encoded protein responds to mTOR (mammalian target of rapamycin) signaling to promote protein synthesis, cell growth, and cell proliferation. Activity of this gene has been associated with human cancer. Alternatively spliced transcript variants have been observed. The use of alternative translation start sites results in isoforms with longer or shorter N-termini which may differ in their subcellular localizations. There are two pseudogenes for this

gene on chromosome 17. [provided by RefSeq, Jan 2013]

Transcript Variant: This variant (1) can initiate translation from two alternate in-frame AUG start codons. The isoform represented in this RefSeq (a, also known as p85 or alpha I) is derived from the first AUG start codon, and is the longest isoform. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the

transcript record were based on transcript alignments.