

Product datasheet for SC324559

RPS15 (NM 001018) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: RPS15 (NM_001018) Human Untagged Clone

Tag:Tag FreeSymbol:RPS15Synonyms:RIG; S15

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-AC (PS100020)E. coli Selection:Ampicillin (100 ug/mL)

Fully Sequenced ORF: >OriGene sequence for NM_001018.3

Restriction Sites: Please inquire **ACCN:** NM_001018

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 001018.3</u>, <u>NP 001009.1</u>

 RefSeq Size:
 531 bp

 RefSeq ORF:
 438 bp

 Locus ID:
 6209

 UniProt ID:
 P62841

 Cytogenetics:
 19p13.3

Domains: Ribosomal_S19

Protein Pathways: Ribosome

Gene Summary: Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and

a large 60S subunit. Together these subunits are composed of 4 RNA species and

approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 40S subunit. The protein belongs to the S19P family of ribosomal

proteins. It is located in the cytoplasm. This gene has been found to be activated in various tumors, such as insulinomas, esophageal cancers, and colon cancers. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome. Alternative splicing results in multiple transcript variants.

[provided by RefSeq, Apr 2015]

Transcript Variant: This variant (2) uses an alternate 5'-terminal exon, which results in a different 5' UTR and use of an alternate start codon compared to variant 1. It encodes isoform

2, which is shorter than and has a distinct N-terminus compared to isoform 1.