



<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_001135699.1</a> , <a href="#">NP_001129171.1</a>
<b>RefSeq Size:</b>	3020 bp
<b>RefSeq ORF:</b>	738 bp
<b>Locus ID:</b>	7534
<b>UniProt ID:</b>	<a href="#">P63104</a>
<b>Cytogenetics:</b>	8q22.3
<b>Protein Pathways:</b>	Cell cycle, Neurotrophin signaling pathway, Oocyte meiosis, Pathogenic Escherichia coli infection
<b>Gene Summary:</b>	<p>This gene product belongs to the 14-3-3 family of proteins which mediate signal transduction by binding to phosphoserine-containing proteins. This highly conserved protein family is found in both plants and mammals, and this protein is 99% identical to the mouse, rat and sheep orthologs. The encoded protein interacts with IRS1 protein, suggesting a role in regulating insulin sensitivity. Several transcript variants that differ in the 5' UTR but that encode the same protein have been identified for this gene. [provided by RefSeq, Oct 2008]</p> <p>Transcript Variant: This variant (3) differs in the 5' UTR compared to variant 2. All six transcripts encode the same protein.</p>