

## Product datasheet for SC324982

### Leukotriene B4 Receptor (LTB4R) (NM\_001143919) Human Untagged Clone

#### Product data:


Product Type:	Expression Plasmids
Product Name:	Leukotriene B4 Receptor (LTB4R) (NM_001143919) Human Untagged Clone
Tag:	Tag Free
Symbol:	LTB4R
Synonyms:	BLT1; BLTR; CMKRL1; GPR16; LTB4R1; LTBR1; P2RY7; P2Y7
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC324982 representing NM_001143919. Blue=Insert sequence Red=Cloning site Green=Tag(s)

```
GCTCGTTTGTAGTGAACCGTCAGAATTTTGTAAACGACTACTATAGGGCGCCGGGAATTCGTCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGAACACTACATCTTCTGCAGCACCCCTCACTAGGTGTAGAGTTCATCTCTCTGCTGGCTATCATC
CTGCTGTGAGTGGCGCTGGCTGTGGGGCTTCCCGCAACAGCTTTGTGGTGTGGAGTATCCTGAAAAGG
ATGCAGAAAGCGCTGTCACTGCCCTGATGGTGTGAACCTGGCCCTGGCCGACCTGGCCGATTGCTC
ACTGCTCCCTTTTCTTCACTTCTGGCCCAAGGCACCTGGAGTTTTGGACTGGCTGGTGGCCGCTG
TGTCACTATGTCTGCGGAGTCAGCATGTACGCCAGCGTCTGCTTATCACGGCCATGAGTCTAGACCGC
TCACTGGCGGTGGCCCGCCCTTTGTGTCCAGAAGCTACGCACCAAGGCATGGCCCGGGGGTGTG
GCAGGCATCTGGGTGTTGTCTTTCTGTGTCGACACCCGTCCTCGCGTACCGCACAGTAGTGCCTGG
AAAACGAACATGAGCCTGTGCTTCCCGCGGTACCCAGCGAAGGGCACCGGGCCTTCCATCTAATCTTC
GAGGCTGTACGGGCTTCTGCTGCCCTTCTGGCTGTGGTGGCCAGCTACTCGGACATAGGGCGTCCGG
CTACAGGCCCAGCGCTTCCGCCGAGCCGCCGACCGGCCGCTGGTGGTGTCTATCATCTGACCTTC
GCCGCTTCTGGCTGCCCTACCAGTGGTGAACCTGGCTGAGGCGGGCCGCGCTGGCCGGCCAGGCC
GCCGGTTAGGGCTCGTGGGAAGCGGCTGAGCCTGGCCCGCAACGTGCTCATCGCACTCGCCTTCTGT
AGCAGCAGCGTGAACCCGTGCTGTACGCGTGCGCCGGCGGGCCTGCTGCGCTCGGCGGGCGTGGCC
TTCGTCGCCAAGCTGCTGGAGGGCACGGGCTCCGAGGCGTCCAGCACGCCCGGGGGCAGCCTGGCC
CAGACCGCTAGGAGCGGCCCGCCGCTCTGGAGCCCGCCCTTCCGAGAGCCTCACTGCCTCCAGCCCT
CTCAAGTTAAACGAACGAAGTAG
AGCGGACCGACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGAT
ATCCTGGATTACAAGGATGACGACGATAAGGTTTAA
```

Restriction Sites: SgfI-RsrII



[View online »](#)

**Plasmid Map:**


**ACCN:** NM\_001143919

**Insert Size:** 1059 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

**OTI Annotation:** This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** [NM\\_001143919.2](#)

**RefSeq Size:** 2723 bp

**RefSeq ORF:** 1059 bp

**Locus ID:** 1241

UniProt ID: [Q15722](#)

Cytogenetics: 14q12

Protein Families: Druggable Genome, GPCR, Transmembrane

Protein Pathways: Neuroactive ligand-receptor interaction

MW: 37.6 kDa

**Gene Summary:** Receptor for extracellular ATP > UTP and ADP. The activity of this receptor is mediated by G proteins which activate a phosphatidylinositol-calcium second messenger system. May be the cardiac P2Y receptor involved in the regulation of cardiac muscle contraction through modulation of L-type calcium currents. Is a receptor for leukotriene B4, a potent chemoattractant involved in inflammation and immune response.[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (2) differs in the 5' UTR, compared to variant 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.