

## **Product datasheet for SC325671**

## VASH2 (NM 001136475) Human Untagged Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** VASH2 (NM\_001136475) Human Untagged Clone

Tag:Tag FreeSymbol:VASH2

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >SC325671 representing NM\_001136475.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

**Restriction Sites:** Sgfl-Mlul




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## Plasmid Map:



**ACCN:** NM\_001136475

**Insert Size:** 756 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.



MW:

## VASH2 (NM\_001136475) Human Untagged Clone - SC325671

**RefSeq:** <u>NM 001136475.2</u>

29.3 kDa

 RefSeq Size:
 4014 bp

 RefSeq ORF:
 756 bp

 Locus ID:
 79805

 UniProt ID:
 Q86V25

 Cytogenetics:
 1q32.3

**Gene Summary:** Tyrosine carboxypeptidase that removes the C-terminal tyrosine residue of alpha-tubulin,

thereby regulating microtubule dynamics and function (PubMed:29146869). Acts as an activator of angiogenesis: expressed in infiltrating mononuclear cells in the sprouting front to

promote angiogenesis (PubMed:19204325).[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (3) contains multiple differences, compared to variant 1, including initiation of translation from a downstream in-frame start codon. The encoded isoform (2) is shorter than isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data because no single transcript was available for the full length of the gene. The extent of this transcript is supported by transcript alignments.