

Product datasheet for SC325722

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Caspase 5 (CASP5) (NM_001136110) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Caspase 5 (CASP5) (NM_001136110) Human Untagged Clone

Tag: Tag Free
Symbol: Caspase 5

Synonyms: ICE(rel)III; ICEREL-III; ICH-3

Vector: <u>pCMV6 series</u>

Fully Sequenced ORF: >NCBI ORF sequence for NM_001136110, the custom clone sequence may differ by one or

more nucleotides

ACCTTGACAAGAGATTTCTACCTCTTTCCTGGCAAT

Restriction Sites: Please inquire ACCN: NM 001136110

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.



Caspase 5 (CASP5) (NM_001136110) Human Untagged Clone - SC325722

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: NM 001136110.1, NP 001129582.1

 RefSeq Size:
 1023 bp

 RefSeq ORF:
 879 bp

 Locus ID:
 838

 UniProt ID:
 P51878

 Cytogenetics:
 11q22.3

Protein Families: Druggable Genome, Protease

Protein Pathways: NOD-like receptor signaling pathway

Gene Summary: This gene encodes a member of the cysteine-aspartic acid protease (caspase) family.

Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. Overexpression of the active form of this enzyme induces apoptosis in fibroblasts. Max, a central component of the Myc/Max/Mad transcription regulation network important for cell growth, differentiation, and apoptosis, is cleaved by this protein; this process requires

Fas-mediated dephosphorylation of Max. The expression of this gene is regulated by

interferon-gamma and lipopolysaccharide. Alternatively spliced transcript variants have been

identified for this gene. [provided by RefSeq, Aug 2010]

Transcript Variant: This variant (c) is lacking two consecutive in-frame coding exons compared to transcript variant a, resulting in a shorter isoform (c) missing a 142 aa protein segment

compared to isoform a.