

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for SC325801

CD299 (CLEC4M) (NM_001144904) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	CD299 (CLEC4M) (NM_001144904) Human Untagged Clone
Tag:	Tag Free
Symbol:	CD299
Synonyms:	CD209L; CD299; DC-SIGN2; DC-SIGNR; DCSIGNR; HP10347; L-SIGN; LSIGN
Vector:	pCMV6 series
Fully Sequenced ORF:	<pre>>NCBI ORF sequence for NM_001144904, the custom clone sequence may differ by one or more nucleotides ATGAGTGACTCCAAGGAACCAAGGGTGCAGCAGCTGGGCCTCCTGGGGGTGCTTGGCCAT GGCGCCCTGGTGCTGCAACTCCTCTCCT</pre>
Restriction Sites:	Please inquire
ACCN:	NM_001144904
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

CD299 (CLEC4M) (NM_001144904) Human Untagged Clone – SC325801		
OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.	
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).	
Reconstitution Method:	 Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C. 	
RefSeq:	<u>NM 001144904.1, NP 001138376.1</u>	
RefSeq Size:	1799 bp	
RefSeq ORF:	1047 bp	
Locus ID:	10332	
Cytogenetics:	19p13.2	
Protein Families:	Druggable Genome, Transmembrane	
Gene Summary:	This gene encodes a C-type lectin that functions in cell adhesion and pathogen recognition. This receptor recognizes a wide range of evolutionarily divergent pathogens with a large impact on public health, including tuberculosis mycobacteria, and viruses including Ebola, hepatitis C, HIV-1, influenza A, West Nile virus and the SARS-CoV acute respiratory syndrome coronavirus. The protein is organized into four distinct domains: a C-terminal carbohydrate recognition domain, a flexible tandem-repeat neck domain of variable length, a transmembrane region and an N-terminal cytoplasmic domain involved in internalization. This gene is closely related in terms of both sequence and function to a neighboring gene, CD209 (Gene ID: 30835), also known as DC-SIGN. The two genes differ in viral recognition and expression patterns, with this gene showing high expression in endothelial cells of the liver, lymph node and placenta. Polymorphisms in the tandem repeat neck domain are associated with resistance to SARS infection. [provided by RefSeq, May 2020] Transcript Variant: This variant (2) lacks an exon and uses an alternative splice site in the coding region and maintains the reading frame. This results in a shorter protein (isoform 2) compared to isoform 1. This isoform (2) has 6.5 repeats in the neck domain.	

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US