

Product datasheet for SC325844

Caspase 5 (CASP5) (NM_001136109) Human Untagged Clone

Product data:


Product Type:	Expression Plasmids
Product Name:	Caspase 5 (CASP5) (NM_001136109) Human Untagged Clone
Tag:	Tag Free
Symbol:	CASP5
Synonyms:	ICE(rel)III; ICEREL-III; ICH-3
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC325844 representing NM_001136109. Blue=Insert sequence Red=Cloning site Green=Tag(s)

```
GCTCGTTTGTAGTGAACCGTCAGAATTTGTAAACGACTCACTATAGGGCGCCGGGAATTCGTGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGGCTGAAGACAACACAAAAAACAAGTTAAGATGTTGGAATACCTGGCAAGATGTTCTTCAT
GGTGTTTTTAATTATTTGGCAAAACACGATGTTCTGACATTGAAGGAAGAGGAAAAGAAAAATATTAT
GATACCAAAATGAAGACAAGGCCCTGATCTTGGTAGACTCTTTGCGAAAGAATCGCGTGGCTCATCAA
ATGTTTACCCAAACACTTCTCAATATGGACCAAAAGATCACCAGTGTAAAACCTTCTCGAAATCGAG
GCTGGACCACCTGAGTCAGCAGAATCTACAAATACTCAAACCTTTGTCCTCGTGAAGAATTCCTGAGA
CTGTGTAATAAATCATGATGAGATCTATCCAATAAAAAAGAGAGAGGACCGCAGACGCCTGGCTCTC
ATCATATGCAATACAAAGTTTGATCACCTGCCTGCAAGGAATGGGGCTCACTATGACATCGTGGGATG
AAAAGGCTGCTTCAAGGCCTGGGCTACACTGTGGTTGACGAAAAGAAATCTCACAGCCAGGGATATGGAG
TCAGTGTGAGGGCATTGCTGCCAGACCAGAGCAAGTCTCTGACAGCACGTTCTTGGTACTCATG
TCTCATGGCATCCTAGAGGGAATCTGCCGAAGTGCATATAAAGAAAAACCGGATGTGCTGCTTTAT
GACACCATCTCCAGATATCAACAACCGCAACTGCCTCAGTCTAAAGGACAAACCAAGGTCATCATT
GTCCAGGCCTGCAGAGGTGAAAAACATGGGAACTCTGGGTCAGAGACTCTCCAGCATCCTTGGCACTC
ATCTCTTCACAGTCATCTGAGAACCTGGAGGCAGATTCTGTTTGAAGATCCACGAGGAGAAGGACTTC
ATTGCTTTCTGTTCTTCAACACCACATAACGTGTCTGGAGAGACCGCACAGGGGCTCCATCTTCATT
ACGGAACATCATCATGCTTCCAGAAATATTCTTGCTGCTGCCACCTAATGGAAATATTTCCGGAAGGTA
CAGAAATCATTTGAAGTCCACAGGCTAAAGCCAGATGCCACCATAGAACGAGCAACCTTGACAAGA
GATTTCTACCTCTTCTGGCAATTGA
ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
```

Restriction Sites: SgfI-MluI



[View online »](#)

Plasmid Map:


ACCN: NM_001136109

Insert Size: 1131 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001136109.1](#)

RefSeq Size: 1275 bp

RefSeq ORF: 1131 bp

Locus ID: 838

UniProt ID: [P51878](#)

Cytogenetics:	11q22.3
Protein Families:	Druggable Genome, Protease
Protein Pathways:	NOD-like receptor signaling pathway
MW:	43.3 kDa
Gene Summary:	<p>This gene encodes a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. Overexpression of the active form of this enzyme induces apoptosis in fibroblasts. Max, a central component of the Myc/Max/Mad transcription regulation network important for cell growth, differentiation, and apoptosis, is cleaved by this protein; this process requires Fas-mediated dephosphorylation of Max. The expression of this gene is regulated by interferon-gamma and lipopolysaccharide. Alternatively spliced transcript variants have been identified for this gene. [provided by RefSeq, Aug 2010]</p> <p>Transcript Variant: This variant (b) is lacking an in-frame coding exon compared to transcript variant a, resulting in a shorter isoform (b) missing a 58 aa protein segment compared to isoform a.</p>