

Product datasheet for SC325929

MDM2 (NM_001145339) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	MDM2 (NM_001145339) Human Untagged Clone
Tag:	Tag Free
Symbol:	MDM2
Synonyms:	ACTFS; hdm2; HDMX; LSKB
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC325929 representing NM_001145339. Blue=Insert sequence Red=Cloning site Green=Tag(s)


```
GCTCGTTTGTGAAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGCCGGGAATTCGTCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGGTGAGGAGCAGGCAAATGTGCAATACCAACATGTCTGTACCTACTGATGGTGTGTAACCACCTCA
CAGATTCAGCTTCGGAACAAGAGACCCTGGTTAGACCAAGCCATTGCTTTTGAAGTTATTAAGTCT
GTTGGTGCACAAAAGACACTTATACTATGAAAGAGGTTCTTTTTATCTTGGCCAGTATATTATGACT
AAACGATTATATGATGAGAAGCAACAACATATTGTATATTGTTCAAATGATCTTCTAGGAGATTTGTTT
GGCGTGCCAAAGCTTCTCTGTGAAAGAGCACAGGAAAATATATACCATGATCTACAGGAACCTGGTAGTA
GTCAATCAGCAGGAAGAAAATTCAGATGAATTATCTGGTGAACGACAAAGAAAACGCCACAAATCTGAT
AGTATTTCCCTTTCTTTGATGAAAGCCTGGCTCTGTGTGTAATAAGGGAGATATGTTGTGAAAGAAGC
AGTAGCAGTGAATCTACAGGGACGCCATCGAATCCGGATCTTGATGCTGGTGAAGTGAACATTCAGGT
GATTGGTTGGATCAGGATTCAGTTTCAGATCAGTTTAGTGTAGAAATTTGAAGTTGAATCTCTCGACTCA
GAAGATTATAGCCTTAGTGAAGAAGGACAAGAAGCTCTCAGATGAAGATGATGAGGTATATCAAGTTACT
GTGTATCAGGCAGGGGAGAGTGATACAGATTCATTTGAAGAAGATCCTGAAATTTCTTAGCTGACTAT
TGGAAATGCACCTCATGCAATGAAATGAATCCCCCCTTCCATCACATTGCAACAGATGTTGGGCCCTT
CGTGAGAATTGGCTTCTGAAGATAAAGGGAAAGATAAAGGGGAAATCTCTGAGAAAGCCAAACTGGAA
AACTCAACACAAGCTGAAGAGGGCTTTGATGTTCTGATTGTAATAAACTATAGTGAATGATTCCAGA
GAGTCATGTGTTGAGGAAAATGATGATAAAATTACACAAGCTTCAATCACAAGAAAGTGAAGACTAT
TCTCAGCCATCAACTTCTAGTAGCATTATTTATAGCAGCCAAGAAGATGTGAAAGAGTTTGAAGGGAA
GAAACCAAGACAAGAAGAGAGTGTGGAATCTAGTTTGGCCCTTAATGCCATTGAACCTTGTGTGATT
TGTCAAGGTCGACCTAAAAATGGTTGCATTGTCCATGGCAAAACAGGACATCTTATGGCCTGCTTTACA
TGTGCAAGAAGCTAAAGAAAAGGAATAAGCCCTGCCAGTATGTAGACAACCAATTCAAATGATTGTG
CTAACTTATTTCCCTAG
ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGCGC
```



[View online »](#)

Restriction Sites: SgfI-MluI

Plasmid Map:



ACCN: NM_001145339

Insert Size: 1329 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001145339.2](#)

RefSeq Size: 7330 bp

RefSeq ORF: 1329 bp

Locus ID: 4193

UniProt ID: [Q00987](#)

Cytogenetics:	12q15
Protein Families:	Druggable Genome, Transcription Factors
Protein Pathways:	Bladder cancer, Cell cycle, Chronic myeloid leukemia, Endocytosis, Glioma, Melanoma, p53 signaling pathway, Pathways in cancer, Prostate cancer, Ubiquitin mediated proteolysis
MW:	50 kDa
Gene Summary:	<p>This gene encodes a nuclear-localized E3 ubiquitin ligase. The encoded protein can promote tumor formation by targeting tumor suppressor proteins, such as p53, for proteasomal degradation. This gene is itself transcriptionally-regulated by p53. Overexpression or amplification of this locus is detected in a variety of different cancers. There is a pseudogene for this gene on chromosome 2. Alternative splicing results in a multitude of transcript variants, many of which may be expressed only in tumor cells. [provided by RefSeq, Jun 2013]</p> <p>Transcript Variant: This variant (2, also known as MDM2g) lacks two coding exons, but maintains the reading frame, compared to variant 1. The encoded isoform (h) is shorter than isoform a. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.</p>