

Product datasheet for **SC325933**

MDM2 (NM_001145337) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	MDM2 (NM_001145337) Human Untagged Clone
Tag:	Tag Free
Symbol:	MDM2
Synonyms:	ACTFS; hdm2; HDMX; LSKB
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC325933 representing NM_001145337. Blue=Insert sequence Red=Cloning site Green=Tag(s)

GCTCGTTTGTAGTAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTG
 GATCCGGTACCGAGGAGATCTGCCGCC**CGCATCGCC**
 ATGTGCAATACCAACATGTCTGTACCTACTGATGGTGCTGAACCACTCACAGATTCAGCTTCGGAA
 CAAGAGACCTGGTTAGACCAAAGCCATTGCTTTTGAAGTTATTAAGTCTGTTGGTGCACAAAAAGAC
 ACTTATACTATGAAAGAGGTTCTTTTTATCTGGCCAGTATATTATGACTAAACGATTATATGATGAG
 AAGCAACAACATATTGTATATTGTTCAATGATCTTCTAGGAGATTTGTTGGCGTGCCAAGCTTCTCT
 GTGAAAGAGCACAGGAAAATATACCATGATCTACAGGAACCTGGTAGTAGTCAATCAGCAGGAATCA
 TCGGACTCAGGTACATCTGTGAGTGAGAACAGGTGTACCTTGAAGGTGGGAGTGATCAAAAGGACCTT
 GTACAAGAGCTTCAGGAAGAGAAACCTTCATCTTCACATTTGGTTTCTAGACCATCTACCTCATCTAGA
 AGGAGAGCAATTAGTGAGACAGAAGAAAATTCAGATGAATTATCTGGTGAACGACAAAAGAAAACGCCAC
 AAATCTGATAGTATTTCCCTTTCTTTGATGAAAGCCTGGCTCTGTGTGAATAAGGGAGATATGTTGT
 GAAAGAAGCAGTAGCAGTGAATCTACAGGGACGCCATCGAATCCGGATCTTGATGCTGGTGTATACAA
 GTTACTGTGTATCAGGCAGGGGAGAGTGATACAGATTCATTTGAAGAAGATCCTGAAATTTCTTAGCT
 GACTATTGGAAATGCACTTCATGCAATGAAATGAATCCCCCTTCCATCACATTGCAACAGATGTTGG
 GCCCTTCGTGAGAATTGGCTTCCTGAAGATAAAGGGAAAGATAAAGGGGAAATCTCTGAGAAAGCCAAA
 CTGGAAAACCTCAACACAAGCTGAAGAGGGCTTTGATGTTCTGATTGTAAAAAACTATAGTGAATGAT
 TCCAGAGAGTCATGTGTTGAGGAAAATGATGATAAAATTACACAAGCTTCACAATCACAAGAAAGTGAA
 GACTATTCTCAGCCATCAACTTCTAGTAGCATTATTTATAGCAGCCAAGAAGATGTGAAAGAGTTTGAA
 AGGGAAGAAACCAAGACAAAGAAGAGAGTGTTGAATCTAGTTTGCCCTTAATGCCATTGAACCTTGT
 GTGATTTGTCAAGGTCGACCTAAAAATGGTTGCATTGTCCATGGCAAAACAGGACATCTTATGGCCTGC
 TTTACATGTGCAAGAAGCTAAAGAAAAGGAATAAGCCCTGCCAGTATGTAGACAACCAATTCAAATG
 ATTGTGCTAACTTATTTCCCT**AG**
ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
 TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC


[View online »](#)

Restriction Sites:	Sgfl-Mlul
ACCN:	NM_001145337
Insert Size:	1335 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001145337.2</u>
RefSeq Size:	7104 bp
RefSeq ORF:	1335 bp
Locus ID:	4193
Cytogenetics:	12q15
Protein Families:	Druggable Genome, Transcription Factors
Protein Pathways:	Bladder cancer, Cell cycle, Chronic myeloid leukemia, Endocytosis, Glioma, Melanoma, p53 signaling pathway, Pathways in cancer, Prostate cancer, Ubiquitin mediated proteolysis
MW:	49.9 kDa

Gene Summary:

This gene encodes a nuclear-localized E3 ubiquitin ligase. The encoded protein can promote tumor formation by targeting tumor suppressor proteins, such as p53, for proteasomal degradation. This gene is itself transcriptionally-regulated by p53. Overexpression or amplification of this locus is detected in a variety of different cancers. There is a pseudogene for this gene on chromosome 2. Alternative splicing results in a multitude of transcript variants, many of which may be expressed only in tumor cells. [provided by RefSeq, Jun 2013] Transcript Variant: This variant (3, also known as P2-MDM2-10) contains multiple differences in the 5' UTR and coding region, compared to variant 1. It uses an alternate promoter and initiates translation at a downstream in-frame start codon. The encoded isoform (g) has a shorter N-terminus and is shorter than isoform a. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.