

## Product datasheet for SC325939

### Caspase 5 (CASP5) (NM\_001136112) Human Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Caspase 5 (CASP5) (NM_001136112) Human Untagged Clone
Tag:	Tag Free
Symbol:	CASP5
Synonyms:	ICE(rel)III; ICEREL-III; ICH-3
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC325939 representing NM_001136112. Blue=Insert sequence Red=Cloning site Green=Tag(s)


```
GCTCGTTTGTGAAACCGTCAGAATTTTGTAAACGACTACTATAGGGCGCCGGGAATTCGTCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGGCTGCTGTGCCAGAGTTGAAGGAGCTTTTATTTCTGATAGAAGACAGTGCCAAAAAAGG
CGTAAGAATTTTGAAGCTATGTTCAAAGGTATCCTTCAGAGTGGATTGGATAACTTCGTGATAAACCAC
ATGCTAAAGAACAACGTGGCTGGACAAACATCTATCCAGACCCTAGTACCTAATACGGATCAAAGTTCG
ACCAGTGTAAAAAAGACAACCACAAAAAAGCAGTTAAGATGTTGGAATACCTGGGCAAGATGTT
CTTCATGGTGTTTTAAATTTTGGCAAAACACGATGTTCTGACATTGAAGGAAGAGGAAAAAGAAAA
TATTATGATACAAAATGAAGACAAGGCCCTGATCTTGGTAGACTCTTTCGAAAAGATCGCGTGGCT
CATCAAATGTTTACCCAAACACTTCTCAATATGGACAAAAGATCACCAGTGTAAAACCTTCTTGCAA
ATCGAGGCTGGACCACCTGAGTCAGCAGAATCTACAAATATACTCAAACCTTGTCCCTCGTGAAGAATTC
CTGAGACTGTGTAATAAATCATGATGAGATCTATCCAATAAAAAAGAGAGAGGACCGCAGACGCTG
GCTCTCATCATATGCAATACAAAGTTTGTACCTGCCTGCAAGGAATGGGGCTCACTATGACATCGTG
GGGATGAAAAGGCTGCTCAAGGCTGGGCTACACTGTGGTTGACGAAAAGAAATCTCACAGCCAGGGAT
ATGGAGTCAGTGTGAGGGCATTGCTGCCAGACCAGAGCACAAGTCTCTGACAGCACGTTCTTGGTA
CTCATGTCTCATGGCATCCTAGAGGGAATCTGCCGAACTGCGCATAAAAAAGAAAAACCGGATGTGCTG
CTTTATGACACCATCTCCAGATATTCAACAACCGCAACTGCCTCAGTCTAAAGGACAAACCAAGGTC
ATCATTGTCCAGGCTGCAGAGGTGAAAAACATGGGGAACCTCTGGGTGAGAGACTCTCCAGCATCCTTG
GCACTCATCTCTCACAGTCATCTGAGAACCTGGAGGCAGATTCTGTTTGAAGATCCACAGGAGAAG
GACTTCATTGCTTCTGTTCTTCAACACCACATAACGTGTCTGGAGAGACCGCACAAGGGGCTCCATC
TTCATTACGGAACCTATCACATGCTTCCAGAAATATTCTTGTGCTGCCACCTAATGAAATATTTTCGG
AAGGTACAGAAATCATTGAAGTCCACAGGCTAAAGCCAGATGCCACCATAGAACGAGCAACCTTG
ACAAGAGATTTCTACCTTTTCTGGCAATTGA
ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGCGC
```



View online »

**Restriction Sites:** SgfI-MluI

**Plasmid Map:**



**ACCN:** NM\_001136112

**Insert Size:** 1344 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

**OTI Annotation:** This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** [NM\\_001136112.1](#)

**RefSeq Size:** 1488 bp

**RefSeq ORF:** 1344 bp

**Locus ID:** 838

**UniProt ID:** [P51878](#)

<b>Cytogenetics:</b>	11q22.3
<b>Protein Families:</b>	Druggable Genome, Protease
<b>Protein Pathways:</b>	NOD-like receptor signaling pathway
<b>MW:</b>	51.2 kDa
<b>Gene Summary:</b>	<p>This gene encodes a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. Overexpression of the active form of this enzyme induces apoptosis in fibroblasts. Max, a central component of the Myc/Max/Mad transcription regulation network important for cell growth, differentiation, and apoptosis, is cleaved by this protein; this process requires Fas-mediated dephosphorylation of Max. The expression of this gene is regulated by interferon-gamma and lipopolysaccharide. Alternatively spliced transcript variants have been identified for this gene. [provided by RefSeq, Aug 2010]</p> <p>Transcript Variant: This variant (f) uses an in-frame, alternate acceptor splice site at an internal coding exon compared to transcript variant a, resulting in a longer isoform (f) containing an additional 13 aa protein segment compared to isoform a.</p>