

Product datasheet for **SC326098**

SGT1 (ECD) (NM_001135753) Human Untagged Clone

Product data:

| | |
|---------------------------|--|
| Product Type: | Expression Plasmids |
| Product Name: | SGT1 (ECD) (NM_001135753) Human Untagged Clone |
| Tag: | Tag Free |
| Symbol: | ECD |
| Synonyms: | GCR2; HSGT1; SGT1 |
| Mammalian Cell Selection: | Neomycin |
| Vector: | pCMV6-Entry (PS100001) |
| E. coli Selection: | Kanamycin (25 ug/mL) |




[View online »](#)

Fully Sequenced ORF: >SC326098 representing NM_001135753.
Blue=Insert sequence Red=Cloning site Green=Tag(s)

```
GCTCGTTTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGGCCGGGAATTCGTGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGGAAGAAACCATGAAGCTTGCTACGATGGAAGACACAGTGGAGTACTGCCTGTTCTGATACCAGAT
GAGTCAAGGGACTCAGATAAACATAAAGAGATTCTTCAGAAGTACATTGAGAGAATAATCACTCGGTTT
GCACCTATGCTGGTCCCTACATCTGGCAGAATCAGCCTTCAATCTTAAATATAAACCTGGGAAAGGA
GGTGTTCCTGCTCATATGTTTGGCGTGACAAAGTTTGGGATAACATTGAGGATGAATGGTTTATTGTT
TATGTAATAAAGCAGATCACAAGGAATTTCCAGAGTTAGTAGCAAGGATTGAAGACAATGATGGTGAA
TTCTTGTTAATAGAAGCTGCTGACTTTCTCCCTAAATGGCTGGATCCTGAAAATAGCACCAATAGGGTA
TTTTTCTGCCATGGGAATTGTGATTATCCCTGCACCAAGAAAATCTGGAGCAGAATCTTGTTACCC
ACCACACCCCAACAATCCACAAGCATTGAATATAATCACAGCACATTCAGAAAAAATACTTGCTTCA
GAATCTATACGAGCTGCTGTGAATAGGCGCATCAGAGGGTACCCAGAAAAAATTCAGGCCTCACTTCAT
CGAGCACACTGCTTCCTCCAGCTGGCATTGTGGCAGTGTAAAGCAGCGCCCAAGATTGGTGGCTGCA
GCAGTCCAGGCATTTTACCTACGAGACCCTATTGACCTGCGAGCTTGTCTGTTTTCAAGACATTCTTG
CCTGAAACACGAATAATGACATCGGCTCATGGATTTGAGATCTTATGCTCCAAATGTAGCCACATTTT
TCTGACTGCAAGAAATCCCTTGTGACTGCCTCACCCTCTGGGCCAGTTTCTTGAAAGTCTGAAAAAG
AATGATTACTTTAAGGGACTGATAGAAGTTTCTGCTCAGTACCGGGAAAGGCTAGAAATGGCAGAGAAT
TACTTCCAGCTCTCAGTAGACTGGCCAGAAAGTTCTCTTGCTATGAGCCCTGGTGAAGAAATCTTAACC
TTATTACAGACAATACCATTTGATATAGAAGACCTTAAGAAAGAAGCAGCTAATCTTCCCCAGAGGAT
GATGACCAGTGGTTAGATCTCTCACCAGATCAGCTGGACCAGCTGCTGCAGGAAGCTGTTGGCAAAAA
GAATCCGAGTCTGTTTTCAAGGAGGAGAAGGAGCAGAACTATGACTTAACTGAAGTCTCAGAGAGCATG
AAAGCTTTCATATCCAAAGTCTCAACCCACAAGGAGCAGAGCTGCCTCGAGAACCCTTCTGAGGCTCCA
ATCACTTTTATGATGAGATTCTTTTCTTAATTTATTTGATAAGATTTTAGGGCCAAGGCCTAATGAGTCA
GATTCTGATGATCTGGATGATGAAGACTTTGAATGTTTAGATAGTATGATGACTTGGACTTTGAAACA
CACGAACCTGGCGAAGAGGCTTCCCTGAAAGGAACACTTGATAATCTCAAGTCATACATGGCCAGATG
GACCAGGAAGTACACACACCTGCATCAGCAAAAGTTTACCCTAGGAACCAAGTGAACCTGTATCC
CAGACTACCGATAACAATTCAGATGAGGAAGATTCTGGTACGGGAGAATCTGTTATGGCACCAGTAGAT
GTAGACCTGAACCTGGTTTCAAATATATTGGAATCCTATAGCTCCAAGCTGGACTGGCAGGACCTGCT
TCCAATCTTTTACAAAGCATGGGAGTGCAGCTGCCTGACAACACCGATCACAGACCAACAAGTAAGCCA
ACAAAAAATTAA
ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
```

Restriction Sites: SgfI-MluI

Plasmid Map:


ACCN: NM_001135753

Insert Size: 1806 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001135753.1](#)

RefSeq Size: 2151 bp

RefSeq ORF: 1806 bp

Locus ID: 11319

UniProt ID: [O95905](#)

Cytogenetics: 10q22.2

| | |
|--------------------------|--|
| Protein Families: | Druggable Genome, Transcription Factors |
| MW: | 67.7 kDa |
| Gene Summary: | <p>Regulator of p53/TP53 stability and function. Inhibits MDM2-mediated degradation of p53/TP53 possibly by cooperating in part with TXNIP (PubMed:16849563, PubMed:23880345). May be involved transcriptional regulation. In vitro has intrinsic transactivation activity enhanced by EP300. May be a transcriptional activator required for the expression of glycolytic genes (PubMed:19919181, PubMed:9928932). Involved in regulation of cell cycle progression. Proposed to disrupt Rb-E2F binding leading to transcriptional activation of E2F proteins (PubMed:19640839). The cell cycle -regulating function may depend on its RUVBL1-mediated association with the R2TP complex (PubMed:26711270). May play a role in regulation of pre-mRNA splicing (PubMed:24722212).[UniProtKB/Swiss-Prot Function]</p> <p>Transcript Variant: This variant includes an additional in-frame coding exon compared to variant 1 resulting in a longer protein isoform (3).</p> |