

Product datasheet for **SC326982**

GM CSF Receptor alpha (CSF2RA) (NM_001161530) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	GM CSF Receptor alpha (CSF2RA) (NM_001161530) Human Untagged Clone
Tag:	Tag Free
Symbol:	CSF2RA
Synonyms:	alphaGMR; CD116; CDw116; CSF2R; CSF2RAX; CSF2RAY; CSF2RX; CSF2RY; GM-CSF-R-alpha; GMCSFR; GMCSFR-alpha; GMR; GMR-alpha; SMDP4
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

Fully Sequenced ORF: >SC326982 representing NM_001161530.
 Blue=Insert sequence Red=Cloning site Green=Tag(s)


```

GCTCGTTTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGGCCGGGAATTCGTGCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGCTTCTCCTGGTGACAAGCCTTCTGCTCTGTGAGTTACCACACCCAGCATTCTCTGATCCCAGAG
AAATCGGATCTGCGAACAGTGGCACCAGCCTCTAGTCTCAATGTGAGTTTGACTCCAGGACGATGAAT
TTAAGCTGGGACTGCCAAGAAAACAACCTTCAGCAAGTGTTCCTAACTGACAAGAAGAACAGAGTC
GTGAACCCAGGCTCAGTAACAACGAATGTTCTGTCACATTTCTGAAATTTGTCTGCATGAAGGAGTC
ACATTTGAGGTTACGTGAATACTAGTCAAAGAGGATTTCAACAGAAACTGCTTTATCCAAATTCAGGA
AGGGAGGGTACCCTGCTCAGAAATTTCTCTGTTTCATCTACAATGCGGATTTAATGAACTGTACCTGG
GCGAGGGTCCGACGGCCCCCGTGACGTCCAGTATTTTTGTACATACGAAACTCAAAGAGAAGGAGG
GAGATCCGGTGCCTTATTACATACAAGACTCAGGAACCCATGTGGGATGCACCTGGATAACCTGTCA
GGATTAACGTCTCGCAATTAATTTCTGGTTAACGGAACCGAGAAATGGCATCCAATTCCTTGAT
TCACTTTTGGACAAAAGAAAATAGAACGATTCAACCCTCCAGCAATGTCACCGTACGTTGCAACACG
ACGCACTGCCTCGTACGGTGAAACAGCCAGGACCTATCAGAAGCTGTGCTACCTGGACTTTCAGTAC
CAGCTGGACGTCCACAGAAAGAATACCCAGCCTGGCACGGAAAACCTACTGATTAATGTTTCTGGTGAT
TTGAAAAATAGATACAATTTCCAAGCTCTGAGCCAGAGCAAAACACAGTGTGAAGATCAGAGCTGCA
GACGTCCGCATCTTGAATTGGAGCTCCTGGAGTGAAGCCATTGAATTTGGGTCTCACTCTGTACCCAG
GCTGGAGTGAATGGACAATCTCGGCTCACTGCAACCTCCGCTCCCAGGCTCAAGCGATTCTCCTGC
CTCAGACTCCCAGTTCTGACGACGGGAACCTCGGCTCTGTGTACATTTATGTGCTCCTAATCGTGGGA
ACCCTTGTCTGTGGCATCGTCTCGGCTTCCTCTTTAAAAGTTTCCTTAGGATACAGCGGCTGTTCCCG
CCAGTCCACAGATCAAAGACAAAATGAATGATAACCATGAGGTGGAAGACGAGATCATCTGGGAGGAA
TTCACCCAGAGGAAGGAAAGGCTACCGGAAGAGGTCTTGACCGTGAAGGAAATTACTGA
ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
  
```

Restriction Sites:

SgfI-MluI

Plasmid Map:



ACCN: NM_001161530

Insert Size: 1305 bp

OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001161530.1</u>
RefSeq Size:	1789 bp
RefSeq ORF:	1305 bp
Locus ID:	1438
UniProt ID:	<u>P15509</u>
Cytogenetics:	X;Y
Protein Families:	Druggable Genome, Secreted Protein, Transmembrane
Protein Pathways:	Cytokine-cytokine receptor interaction, Hematopoietic cell lineage, Jak-STAT signaling pathway, Pathways in cancer
MW:	49.9 kDa
Gene Summary:	<p>The protein encoded by this gene is the alpha subunit of the heterodimeric receptor for colony stimulating factor 2, a cytokine which controls the production, differentiation, and function of granulocytes and macrophages. The encoded protein is a member of the cytokine family of receptors. This gene is found in the pseudoautosomal region (PAR) of the X and Y chromosomes. Multiple transcript variants encoding different isoforms have been found for this gene, with some of the isoforms being membrane-bound and others being soluble. [provided by RefSeq, Jul 2008]</p> <p>Transcript Variant: This variant (8) has an additional in-frame coding exon in the 3' region compared to variant 1. The resulting isoform (f) is the longest, and has identical N- and C-termini compared to isoform a.</p>