

Product datasheet for SC327351

DAOA (NM 001161814) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: DAOA (NM_001161814) Human Untagged Clone

Tag: Tag Free Symbol: DAOA

Synonyms: LG72; SG72

Mammalian Cell None

Selection:

Vector: pCMV6-XL5

E. coli Selection: Ampicillin (100 ug/mL)

Fully Sequenced ORF: >NCBI ORF sequence for NM_001161814, the custom clone sequence may differ by one or

more nucleotides

ATGGCACAGAGGCATTTACAGAGATCATTATGTCCTTGGGTCTCTTACCTTCCTCAGCCC
TATGCAGAGCTTGAAGAAGTAAGCAGCCATGTTGGAAAAGTCTTCATGGCAAGAAACTAT
GAGTTCCTTGCCTATGAGGCCTCTAAGGACCGCAGGCAGCCTCTAGAACGAATGTGGACC
TGCAACTACAACCAGCAAAAAAGACCAGTCATGCAACCACAAGGAAATAACTTCTACCAAA

GCTGAA

Restriction Sites: Please inquire **ACCN:** NM_001161814

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: NM 001161814.1, NP 001155286.1

RefSeq Size: 1009 bp RefSeq ORF: 249 bp Locus ID: 267012 **UniProt ID:** P59103 Cytogenetics: 13q34

Protein Families: Druggable Genome

Gene Summary: This gene encodes a protein that may function as an activator of D-amino acid oxidase, which

> degrades the gliotransmitter D-serine, a potent activator of N-methyl-D-aspartate (NMDA) type glutamate receptors. Studies also suggest that one encoded isoform may play a role in mitochondrial function and dendritic arborization. Polymorphisms in this gene have been implicated in susceptibility to schizophrenia and bipolar affective disorder. Alternatively spliced transcript variants encoding different isoforms have been identified.[provided by

RefSeg, Mar 2011]

Transcript Variant: This variant (3) has an alternate 5' exon and an additional segment in the 5' coding region, as compared to variant 1, which results in a downstream AUG start codon. The resulting isoform (3) is shorter and has a different N-terminus, as compared to isoform 1. CCDS Note: This CCDS ID represents the protein described in PMIDs: 12364586 and 14966479. This transcript is supported by AY170469.2. It should be noted this transcript is predicted to undergo nonsense-mediated mRNA decay (NMD). However, the protein is represented because it was detected endogenously in PMID: 18544534. It is likely that the majority of transcripts representing this variant will undergo NMD, while some low level of NMD escape may allow for the expression of this protein. It is likely that the majority of transcripts representing this variant will undergo NMD, while some low level of NMD escape