

Product datasheet for SC327437

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NT5C3 (NT5C3A) (NM_001166118) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: NT5C3 (NT5C3A) (NM_001166118) Human Untagged Clone

Tag: Tag Free Symbol: NT5C3A

Synonyms: cN-III; hUMP1; NT5C3; P5'N-1; P5N-1; p36; PN-I; POMP; PSN1; UMPH; UMPH1

Mammalian Cell

Selection:

None

Vector: pCMV6-XL5

E. coli Selection: Ampicillin (100 ug/mL)

Fully Sequenced ORF: >NCBI ORF sequence for NM_001166118, the custom clone sequence may differ by one or

more nucleotides

TTACAGAAGATTCTA

Restriction Sites: Please inquire **ACCN:** NM 001166118

Insert Size: 1932 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).



NT5C3 (NT5C3A) (NM_001166118) Human Untagged Clone - SC327437

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001166118.1</u>, <u>NP 001159590.1</u>

 RefSeq Size:
 1932 bp

 RefSeq ORF:
 1932 bp

 Locus ID:
 51251

 UniProt ID:
 Q9H0P0

 Cytogenetics:
 7p14.3

Protein Families: Transmembrane

Protein Pathways: Metabolic pathways, Nicotinate and nicotinamide metabolism, Purine metabolism, Pyrimidine

metabolism

Gene Summary: This gene encodes a member of the 5'-nucleotidase family of enzymes that catalyze the

dephosphorylation of nucleoside 5'-monophosphates. The encoded protein is the type 1 isozyme of pyrimidine 5' nucleotidase and catalyzes the dephosphorylation of pyrimidine 5' monophosphates. Mutations in this gene are a cause of hemolytic anemia due to uridine 5-prime monophosphate hydrolase deficiency. Alternatively spliced transcript variants

encoding multiple isoforms have been observed for this gene, and pseudogenes of this gene are located on the long arm of chromosomes 3 and 4. [provided by RefSeq, Mar 2012] Transcript Variant: This variant (4) differs in the 5' UTR, includes two alternate internal exons,

and initiates translation at a downstream, in-frame start codon, compared to variant 1. The encoded isoform (3) has a shorter N-terminus, compared to isoform 1. Variants 4 and 5 both

encode the same isoform (3).