

Product datasheet for **SC327495**

PRKAR1B (NM_001164759) Human Untagged Clone

Product data:

| | |
|---------------------------|--|
| Product Type: | Expression Plasmids |
| Product Name: | PRKAR1B (NM_001164759) Human Untagged Clone |
| Tag: | Tag Free |
| Symbol: | PRKAR1B |
| Synonyms: | PRKAR1 |
| Mammalian Cell Selection: | None |
| Vector: | <u>pCMV6-XL5</u> |
| E. coli Selection: | Ampicillin (100 ug/mL) |
| Fully Sequenced ORF: | >NCBI ORF sequence for NM_001164759, the custom clone sequence may differ by one or more nucleotides ATGGCCTCCCCGCCCGCTGCCCTCGGAGGAGGACGAGAGCCTGAAGGGCTGTGAGCTG TACGTGCAGCTGCACGGATCCAGCAGGTCTCAAAGACTGTATCGTCCACCTCTGCATC TCCAAGCCGAACGCCCATGAAGTTCCTCCGGGAGCACTTCGAGAAGCTGGAGAAGGAA GAAAACAGGCAGATTTTGGCGCGCAAAAGTCAAACCTCACAGTCGGACTCCCATGATGAG GAGGTGTCGCCACCCCGAACCCTGTGGTGAAGGCCCGCGCGGAGGAGGCGTG AGTGCCGAGGTGTACACCGAGGAGGACGCCGTGCTACGTGAGGAAGGTGATCCCAAG GACTACAAAACCATGACTGCGCTGGCCAAGGCCATCTCCAAGAAGCTGCTTTCGCTCAC CTGGATGACAACGAGAGGAGTGACATATTCGATGCCATGTTCCCTGTCACTCACATCGCT GGGGAGACTGTTATACAGCAAGGAATGAAGGAGACAACCTCTATGTCGTTGATCAAGGG GAAGTGGATGTGTACGTGAACGGAGAGTGGGTGACCAACATCAGCGAGGGAGGCAGCTTC GGGGAGCTGGCGCTCATCTACGGCACCCCGAGGCTGCGACCGTAAAAGCCAAGACGGAC CTCAAGCTCTGGGGGATCGACCGGGACAGCTACCGGCGCATCCTTATGGGCAGCACGCTG AGGAAACGCAAGATGTACGAGGAGTTCTCAGCAAGGTCTCCATCCTAGAGTCCCTGGAG AAGTGGGAGCGTCTGACCGTGGCGGATGCGCTGGAGCCCGTCCAGTTTGAAGATGGAGAG AAAATTGTGGTCCAGGGAGAGCCTGGGGACGACTTTTACATCATCACGGAGGGCACCCGG TCCGTGCTGCAGCGCCGGTCCCCAATGAGGAGTACGTGGAGGTGGGGCGCCTGGGACCC TCTGACTACTTCGGGGAGATTGCACTGCTGCTGAACCGGCCCGGGCGGCACTGTGCTG GCCCGGGGCCCTCAAGTGTGTGAAGCTGGACCGCCCGCTTCGAGCGTGTGCTGGGG CCCTGCTCTGAGATCCTCAAGAGAACATTACGCGTTACAACAGCTTCATCTCCCTCACC GTC |
| Restriction Sites: | Please inquire |
| ACCN: | NM_001164759 |



[View online »](#)

| | |
|-------------------------------|--|
| OTI Disclaimer: | Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP). |
| OTI Annotation: | This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA. |
| Components: | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water). |
| Reconstitution Method: | <ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C. |
| RefSeq: | <u>NM_001164759.1</u> , <u>NP_001158231.1</u> |
| RefSeq Size: | 2423 bp |
| RefSeq ORF: | 1146 bp |
| Locus ID: | 5575 |
| UniProt ID: | <u>P31321</u> |
| Cytogenetics: | 7p22.3 |
| Protein Families: | Druggable Genome |
| Protein Pathways: | Apoptosis, Insulin signaling pathway |
| Gene Summary: | <p>The protein encoded by this gene is a regulatory subunit of cyclic AMP-dependent protein kinase A (PKA), which is involved in the signaling pathway of the second messenger cAMP. Two regulatory and two catalytic subunits form the PKA holoenzyme, disbands after cAMP binding. The holoenzyme is involved in many cellular events, including ion transport, metabolism, and transcription. Several transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Aug 2015]</p> <p>Transcript Variant: This variant (4) differs in the 5' UTR compared to variant 1. All of the variants encode the same protein. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.</p> |