

Product datasheet for **SC327888**

PCTAIRE1 (CDK16) (NM_033018) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	PCTAIRE1 (CDK16) (NM_033018) Human Untagged Clone
Tag:	Tag Free
Symbol:	CDK16
Synonyms:	PCTAIRE; PCTAIRE1; PCTGAIRE; PCTK1
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)


Fully Sequenced ORF: >SC327888 representing NM_033018.
 Blue=Insert sequence Red=Cloning site Green=Tag(s)

```

GCTCGTTTAGTGAACCGTCAGAATTTTGTAAATACGACTCACTATAGGGCGGCCGGGAATTCGTGACTG
GATCCGGTACCGAGGAGATCTGCCGCCACGATCGCC
ATGCAGTCCGAGATCGCCATGGATCGGATGAAGAAGATCAAACCGCAGCTGTCAATGACACTCCGAGGT
GGCCGAGGCATAGACAAGACCAATGGTGCCCTGAGCAGATAGGCTGGATGAGAGTGGTGGTGGC
GGCAGTGACCCTGGAGAGGCCCCACACGTGCTGCTCCTGGGAACTTCGTTCTGCACGGGGCCCACTC
AGCTCTGCACCAGAGATTGTGCACGAGGACTTGAAGATGGGGTCTGATGGGGAGAGTGACCAGGCTTCA
GCCACGTCCTCGGATGAGGTGCAGTCTCCAGTGAGAGTGCATGCGCAACCATCCCCACGCAAGATC
TCCACTGAGGACATCAACAAGCGCCTATCACTACCAGCTGACATCCGGCTGCCTGAGGGCTACCTGGAG
AAGCTGACCCTCAATAGCCCCATCTTTGACAAGCCCTCAGCCGCCCTCCGTCGTGTCAGCCTATCT
GAGATTGGCTTTGGAACTGGAGACCTACATTAAGCTGGACAACTGGGCGAGGGTACCTATGCCACC
GTCTACAAAGGCAAAAGCAAGCTCACAGACAACCTTGTGGCACTCAAGGAGATCAGACTGGAACATGAA
GAGGGGGCACCCCTGCACCGCCATCCGGGAAGTGTCCCTGCTCAAGGACCTCAAACACGCCAACATCGTT
ACGCTACATGACATTATCCACACGGAGAAGTCCCTCACCCTTGTCTTTGAGTACCTGGACAAGGACCTG
AAGCAGTACCTGGATGACTGTGGGAACATCATCAACATGCACAACGTGAAACTGTTCTGTTCACGCTG
CTCCGTGGCCTGGCCTACTGCCACCGGCAGAAGGTGTACACCGAGACCTCAAGCCCCAGAACCTGCTC
ATCAACGAGAGGGGAGAGCTCAAGCTGGCTGACTTTGGCCTGGCCCGAGCCAAGTCAATCCCAACAAAG
ACATACTCCAATGAGGTGGTGACACTGTGGTACCGGCCCTGACATCCTGCTTGGGTCCACGGACTAC
TCCACTCAGATTGACATGTGGGGTGTGGCTGCATCTTCTATGAGATGGCCACAGGCCGTCCTCTTT
CCGGGCTCCACGGTGGAGAACAGCTACACTTCATCTTCCGTATCTTAGGAACCCCAACTGAGGAGAGC
TGCCACAGGCATCCTGTCCAACGAGGAGTTCAAGACATACAACTACCCCAAGTACCGAGCCGAGCCCTT
TTGAGCCACGCACCCCGACTTGATAGCGACGGGGCCGACCTCCTACCAAGCTGTTGCAGTTTGAGGGT
CGAAATCGGATCTCCGCAGAGGATGCCATGAAACATCCATTCTTCTCAGTCTGGGGGAGCGGATCCAC
AACTTCTGACACTACTTCCATATTTGACTAAAGGAGATTAGCTACAAAAGGAGGCCAGCCTTCGG
TCTTCGTCGATGCCTGACTCAGGCAGGCCAGCTTTCGCGTGGTGGACACCGAGTTCTAA
ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
  
```

Restriction Sites: SgfI-MluI

Plasmid Map:



ACCN: NM_033018

Insert Size:	1509 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_033018.3
RefSeq Size:	3280 bp
RefSeq ORF:	1509 bp
Locus ID:	5127
UniProt ID:	Q00536
Cytogenetics:	Xp11.3
Domains:	pkinase, TyrKc, S_TKc
Protein Families:	Druggable Genome, Protein Kinase
MW:	56.4 kDa
Gene Summary:	<p>The protein encoded by this gene belongs to the cdc2/cdkx subfamily of the ser/thr family of protein kinases. It may play a role in signal transduction cascades in terminally differentiated cells; in exocytosis; and in transport of secretory cargo from the endoplasmic reticulum. This gene is thought to escape X inactivation. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Dec 2009]</p> <p>Transcript Variant: This variant (2) uses an alternate exon in the 5' UTR and 5' coding region, compared to variant 1. It encodes isoform 2, which has a longer and distinct N-terminus, compared to isoform 1.</p>