

## Product datasheet for **SC328167**

### DAP12 (TYROBP) (NM\_001173515) Human Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	DAP12 (TYROBP) (NM_001173515) Human Untagged Clone
Tag:	Tag Free
Symbol:	TYROBP
Synonyms:	DAP12; KARAP; PLOSL; PLOSL1
Mammalian Cell Selection:	None
Vector:	<u>pCMV6-XL5</u>
E. coli Selection:	Ampicillin (100 ug/mL)
Fully Sequenced ORF:	>NCBI ORF sequence for NM_001173515, the custom clone sequence may differ by one or more nucleotides ATGGGGGGACTTGAACCCTGCAGCAGGCTCCTGCTCCTGCCTCCTCCTGCTGGCTGTAAGT GATTGCAGTTGCTCTACGGTGAGCCCGGCGTGCTGGCAGGGATCGTGATGGGAGACCTG GTGCTGACAGTGCTCATTGCCCTGGCCGTACTTCTGCGCCGCTGGTCCCTCGGGG CGAGGGGCTGCGGAGGCGACCCGAAACAGCGTATCACTGAGACCGAGTCGCCTTATCAG GAGCTCCAGGGTCAGAGGTCGGATGTCTACAGCGACCTCAACACACAGAGGCCGTATTAC AAATGA
Restriction Sites:	Please inquire
ACCN:	NM_001173515
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



[View online »](#)

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** [NM\\_001173515.1](#), [NP\\_001166986.1](#)

**RefSeq Size:** 572 bp

**RefSeq ORF:** 306 bp

**Locus ID:** 7305

**Cytogenetics:** 19q13.12

**Protein Families:** Druggable Genome, Transmembrane

**Protein Pathways:** Natural killer cell mediated cytotoxicity

**Gene Summary:** This gene encodes a transmembrane signaling polypeptide which contains an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain. The encoded protein may associate with the killer-cell inhibitory receptor (KIR) family of membrane glycoproteins and may act as an activating signal transduction element. This protein may bind zeta-chain (TCR) associated protein kinase 70kDa (ZAP-70) and spleen tyrosine kinase (SYK) and play a role in signal transduction, bone modeling, brain myelination, and inflammation. Mutations within this gene have been associated with polycystic lipomembranous osteodysplasia with sclerosing leukoencephalopathy (PLOSL), also known as Nasu-Hakola disease. Its putative receptor, triggering receptor expressed on myeloid cells 2 (TREM2), also causes PLOSL. Multiple alternative transcript variants encoding distinct isoforms have been identified for this gene. [provided by RefSeq, Mar 2010]  
Transcript Variant: This variant (4) lacks an alternate in-frame exon and uses an alternate in-frame splice site in the coding region, compared to variant 1. The resulting isoform (4) is shorter compared to isoform 1.