

# Product datasheet for SC328300

## VEGFA (NM\_001171626) Human Untagged Clone

### **Product data:**

Product Type:	Expression Plasmids
Product Name:	VEGFA (NM_001171626) Human Untagged Clone
Tag:	Tag Free
Symbol:	VEGFA
Synonyms:	MVCD1; VEGF; VPF
Mammalian Cell Selection:	None
Vector:	pCMV6-XL4
E. coli Selection:	Ampicillin (100 ug/mL)

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### Fully Sequenced ORF:

>OriGene sequence for NM\_001171626 edited GGCGACTCGGCGCTCGGAAGCCGGGCTCATGGACGGGTGAGGCGGCGGTGTGCGCAGACA GTGCTCCAGCCGCGCGCGCCCCAGGCCCTGGCCCGGGCCCGGGGAGGAAGAG TAGCTCGCCGAGGCGCCGAGGAGAGCGGGCCGCCCCACAGCCCGAGCCGGAGAGGGAGCG CGAGCCGCCGGCCCCGGTCGGGCCTCCGAAACCATGAACTTTCTGCTGTCTTGGGTGC ATTGGAGCCTTGCCTGCTGCTCTACCTCCACCATGCCAAGTGGTCCCAGGCTGCACCCA TGGCAGAAGGAGGAGGGCAGAATCATCACGAAGTGGTGAAGTTCATGGATGTCTATCAGC GCAGCTACTGCCATCCAATCGAAACCCTGGTGGACATCTTCCAGGAGTACCCTGATGAGA TCGAGTACATCTTCAAGCCATCCTGTGTGCCCCTGATGCGATGCGGGGGGCTGCTGCAATG ACGAGGGCCTGGAGTGTGTGCCCACTGAGGAGTCCAACATCACCATGCAGATTATGCGGA TCAAACCTCACCAAGGCCAGCACATAGGAGAGATGAGCTTCCTACAGCACAACAAATGTG AATGCAGACCAAAGAAAGATAGAGCAAGACAAGAAAAATCCCTGTGGGCCTTGCTCAGAGC GGAGAAAGCATTTGTTTGTACAAGATCCGCAGACGTGTAAATGTTCCTGCAAAAACACAG CGAGGCGGTGAGCCGGGCAGGAGGAAGGAGCCTCCCTCAGGGTTTCGGGAACCAGATCTC TCACCAGGAAAGACTGATACAGAACGATCGATACAGAAACCACGCTGCCGCCACCACACC CCCAGCACGGTCCCTCTTGGAATTGGATTCGCCATTTTATTTTTCTTGCTGCTAAATCAC ATATAAAATATATATATTCTTTTTTTTTTTAAATTAACAGTGCTAATGTTATTGGTGTCTTCAC TGGATGTATTTGACTGCTGTGGACTTGAGTTGGGAGGGGAATGTTCCCACTCAGATCCTG ACAGGGAAGAGGAGGAGATGAGAGACTCTGGCATGATCTTTTTTTGTCCCACTTGGTGG GGCCAGGGTCCTCTCCCCTGCCCAGGAATGTGCAAGGCCAGGGCATGGGGGCAAATATGA CCCAGTTTTGGGAACACCGACAAACCCAGCCCTGGCGCTGAGCCTCTCTACCCCAGGTCA GACGGACAGAAAGACAGATCACAGGTACAGGGATGAGGACACCGGCTCTGACCAGGAGTT TGGGGAGCTTCAGGACATTGCTGTGCTTTGGGGATTCCCTCCACATGCTGCACGCGCATC TCGCCCCAGGGGCACTGCCTGGAAGATTCAGGAGCCTGGGCGGCCTTCGCTTACTCTCA CCTGCTTCTGAGTTGCCCAGGAGACCACTGGCAGATGTCCCGGCGAAGAGAAGAGAAGACACA TTGTTGGAAGAAGCAGCCCATGACAGCTCCCCTTCCTGGGACTCGCCCTCATCCTCTCC TGCTCCCCTTCCTGGGGTGCAGCCTAAAAGGACCTATGTCCTCACACCATTGAAACCACT AGTTCTGTCCCCCCAGGAGACCTGGTTGTGTGTGTGTGAGTGGTTGACCTTCCTCCATCC CCTGGTCCTTCCCTTCCCGAGGCACAGAGAGAGAGAGGGCAGGATCCACGTGCCCAT TGTGGAGGCAGAGAAAAGAGAAAGTGTTTTATATACGGTACTTATTTAATATCCCTTTTT GTCACTAGCTTATCTTGAACAGATATTTAATTTTGCTAACACTCAGCTCTGCCCTCCCCG ATCCCCTGGCTCCCCAGCACACATTCCTTTGAAATAAGGTTTCAATATACATCTACATAC TATGTGATTCTGATAAAATAGACATTGCTATTCTGTTTTTTATATGTAAAAAACAAAACAA GAAAAAATAGAGAATTCTACATACTAAATCTCTCTCCTTTTTTAATTTTAATATTTGTTA TCATTTATTGGTGCTACTGTTTATCCGTAATAATTGTGGGGAAAAGATATTAACAT 

Restriction Sites: ACCN: Insert Size:

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Please inquire

NM\_001171626

2770 bp

#### 

OTI Disclaimer:	Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <u>custsupport@origene.com</u> or by calling 301.340.3188 option 3 for pricing and delivery.
	reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	The ORF of this clone has been fully sequenced and found to be a perfect match to NM_001171626.1.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol> <li>Centrifuge at 5,000xg for 5min.</li> <li>Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>Close the tube and incubate for 10 minutes at room temperature.</li> <li>Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	<u>NM 001171626.1, NP 001165097.1</u>
RefSeq Size:	3554 bp
RefSeq ORF:	576 bp
Locus ID:	7422
UniProt ID:	<u>P15692</u>
Cytogenetics:	6p21.1
Protein Families:	Druggable Genome, Secreted Protein
Protein Pathways:	Bladder cancer, Cytokine-cytokine receptor interaction, Focal adhesion, mTOR signaling pathway, Pancreatic cancer, Pathways in cancer, Renal cell carcinoma, VEGF signaling pathway

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### VEGFA (NM\_001171626) Human Untagged Clone – SC328300

Gene Summary:

This gene is a member of the PDGF/VEGF growth factor family. It encodes a heparin-binding protein, which exists as a disulfide-linked homodimer. This growth factor induces proliferation and migration of vascular endothelial cells, and is essential for both physiological and pathological angiogenesis. Disruption of this gene in mice resulted in abnormal embryonic blood vessel formation. This gene is upregulated in many known tumors and its expression is correlated with tumor stage and progression. Elevated levels of this protein are found in patients with POEMS syndrome, also known as Crow-Fukase syndrome. Allelic variants of this gene have been associated with microvascular complications of diabetes 1 (MVCD1) and atherosclerosis. Alternatively spliced transcript variants encoding different isoforms have been described. There is also evidence for alternative translation initiation from upstream non-AUG (CUG) codons resulting in additional isoforms. A recent study showed that a C-terminally extended isoform is produced by use of an alternative inframe translation termination codon via a stop codon readthrough mechanism, and that this isoform is antiangiogenic. Expression of some isoforms derived from the AUG start codon is regulated by a small upstream open reading frame, which is located within an internal ribosome entry site. The levels of VEGF are increased during infection with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), thus promoting inflammation by facilitating recruitment of inflammatory cells, and by increasing the level of angiopoietin II (Ang II), one of two products of the SARS-CoV-2 binding target, angiotensin-converting enzyme 2 (ACE2). In turn, Ang II facilitates the elevation of VEGF, thus forming a vicious cycle in the release of inflammatory cytokines. [provided by RefSeq, Jun 2020] Transcript Variant: This variant (4) lacks an alternate in-frame exon in the 3' coding region compared to variant 1. It can initiate translation from four non-AUG (CUG) sites, and also from an in-frame downstream AUG. This RefSeq represents isoform VEGF-A (also referred to as VEGF165 or VEGF-A165), which is derived from the AUG start codon, and is shorter than isoform a. This variant also encodes a C-terminally extended isoform (VEGF-Ax) that results from the use of an alternative in-frame, downstream translation termination codon (UGA). Isoform VEGF-Ax is represented in NM\_001317010.1. CCDS Note: This CCDS, which is supported by the mRNA BC058855.1, represents a VEGFA isoform derived from an AUG start codon. Alternative translation initiation from multiple upstream non-AUG(CUG) codons is also possible for this gene, as indicated in PMIDs:11352659, 11563986 and 11731620. The longest isoform encoded by this variant, which is derived from the 5'-most CUG start codon, is represented by CCDS 34458.1.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US