

## Product datasheet for **SC328529**

### BCAT1 (NM\_001178092) Human Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	BCAT1 (NM_001178092) Human Untagged Clone
Tag:	Tag Free
Symbol:	BCAT1
Synonyms:	BCATC; BCT1; ECA39; MECA39; PNAS121; PP18
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC328529 representing NM_001178092. Blue=Insert sequence Red=Cloning site Green=Tag(s)

GCTCGTTTGTAGTGAACCGTCAGAATTTTGTAAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTG  
 GATCCGGTACCGAGGAGATCTGCCGCC**CGATCGCC**  
 ATGAAGGCTAAAGACCTAATAGTCACACCAGCTACCATTTTAAAGGAAAAACCAGACCCAATAATCTG  
 GTTTTGGAACTGTGTTACGGATCATATGCTGACGGTGGAGTGGTCCTCAGAGTTTGGATGGGAGAAA  
 CCTCATATCAAGCCTCTCAGAACCTGTCATTGCACCCTGGCTCATCAGCTTTCAGCTATGCAGTGGAA  
 GTATTTGACAAAGAAGAGCTCTTAGAGTGTATTCAACAGCTTGTGAAATTGGATCAAGAATGGGTCCCA  
 TATTCAACATCTGCTAGTCTGTATATTCGTCCTACATTATTGGAAGTGAAGCTTCTCTGGAGTCAAG  
 AAGCCTACCAAAGCCCTGCTCTTTGTACTCTTGAGCCAGTGGGACCTTATTTTCAAGTGAACCTTT  
 AATCCAGTGTCCCTGTGGGCAATCCCAAGTATGTAAGAGCCTGGAAAGGTGGAAGTGGGGACTGCAAG  
 ATGGGAGGGAATTACGGCTCATCTCTTTTGCCTAATGTGAAGCAGTAGATAATGGGTGTCAGCAGGTC  
 CTGTGGCTCTATGGAGAGGACCATCAGATCACTGAAGTGGGAAGTATGAATCTTTTCTTTACTGGATA  
 AATGAAGATGGAGAAGAAGAACTGGCAACTCCTCCACTAGATGGCATCATTCTCCAGGAGTGACAAGG  
 CGGTGCATTCTGGACCTGGCACATCAGTGGGTGAATTTAAGGTGTCAGAGAGATACCTCACCATGGAT  
 GACTTGACAACAGCCCTGGAGGGGAACAGAGTGAGAGAGATGTTTGGCTCTGGTACAGCCTGTGTTGTT  
 TGCCAGTTTCTGATATACTGTACAAAGCGGAGACAATACACATTCCAATATGGAGAATGGTCCTAAG  
 CTGGCAAGCCGCATCTTGAGCAAATTAAGTATCCAGTATGGAAGAGAAGAGAGCGACTGGACAATT  
 GTGCTATCT**GA**  
**ACGCGT**ACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGAT  
 TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

Restriction Sites:	SgfI-MluI
ACCN:	NM_001178092
Insert Size:	978 bp


[View online »](#)

<b>OTI Disclaimer:</b>	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
<b>OTI Annotation:</b>	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"> <li>1. Centrifuge at 5,000xg for 5min.</li> <li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>3. Close the tube and incubate for 10 minutes at room temperature.</li> <li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
<b>RefSeq:</b>	<a href="#">NM_001178092.1</a>
<b>RefSeq Size:</b>	9499 bp
<b>RefSeq ORF:</b>	978 bp
<b>Locus ID:</b>	586
<b>UniProt ID:</b>	<a href="#">P54687</a>
<b>Cytogenetics:</b>	12p12.1
<b>Protein Families:</b>	Druggable Genome
<b>Protein Pathways:</b>	Metabolic pathways, Pantothenate and CoA biosynthesis, Valine, leucine and isoleucine biosynthesis, Valine, leucine and isoleucine degradation
<b>MW:</b>	36.3 kDa
<b>Gene Summary:</b>	<p>This gene encodes the cytosolic form of the enzyme branched-chain amino acid transaminase. This enzyme catalyzes the reversible transamination of branched-chain alpha-keto acids to branched-chain L-amino acids essential for cell growth. Two different clinical disorders have been attributed to a defect of branched-chain amino acid transamination: hypervalinemia and hyperleucine-isoleucinemia. As there is also a gene encoding a mitochondrial form of this enzyme, mutations in either gene may contribute to these disorders. Alternatively spliced transcript variants have been described. [provided by RefSeq, May 2010]</p> <p>Transcript Variant: This variant (3) lacks two in-frame exons in the 5' coding region, compared to variant 1. This results in a shorter protein (isoform 3), compared to isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.</p>