

Product datasheet for **SC328647**

MCG10 (PCBP4) (NM_001174100) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	MCG10 (PCBP4) (NM_001174100) Human Untagged Clone
Tag:	Tag Free
Symbol:	PCBP4
Synonyms:	CBP; LIP4; MCG10
Mammalian Cell Selection:	None
Vector:	<u>pCMV6-XL5</u>
E. coli Selection:	Ampicillin (100 ug/mL)
Fully Sequenced ORF:	>NCBI ORF sequence for NM_001174100, the custom clone sequence may differ by one or more nucleotides

```

ATGAGCGGCTCGGACGGGGGACTGGAGGAGGAGCCAGAGCTCAGCATCACCTCAGCTG
CGGATGCTGATGCACGGGAAGGAAGTGGGCAGCATCATCGGGAAGAAGGGCGAGACTGTA
AAGCGAATCCGGGAGCAGAGCAGTGCCCGGATCACCATCTCCGAGGGCTCCTGCCCTGAA
CGCATCACCACTCACCGGGTCTACAGCAGCTGTCTTCCATGCAGTCTCCATGATTGCT
TTCAAACCTGGATGAGGACCTTTGTGCTGCTCCTGCAAATGGTGGAAATGTCTCCAGGCCT
CCAGTGACCTGCGCCTTGTTCATCCCTGCCAGTCAGTGTGGTCACTGATTGGGAAGGCT
GGCACCAAGATCAAGGAGATCCGAGAGACTACGGGTGCCAGGTACAGGTGGCAGGGGAC
CTGCTCCCAACTCCACAGAGCGAGCTGTTACGGTATCTGGGGTGCCTGATGCCATCATC
CTGTGTGTGCGCCAGATCTGCGCTGTTATCCTGGAGTCCCACCCAAAGGAGCCACTATC
CCCTACCATCCGAGCCTCTCCCTAGGTAAGTCTTCTCTCTGCCAACCAGGGCTTCTCT
GTCCAGGGTCAGTATGGGGCTGTGACCCAGCTGAGGTACCAAGCTCCAGCAGCTCTCA
AGCCATGCGGTCCCCTTTGCCACACCCAGCGTGGTCCAGGACTGGATCCCAGCACACAG
ACCAGCTCACAGGAGTCTTGGTTCCCAACGATTTGATTGGTGTGTGATCGGGGCCAG
GGCAGCAAGATCAGCGAGATCCGGCAGATGTCAGGGGCACATATCAAGATCGGGAACCA
GCAGAGGGCGCTGGGGAGCGGCATGTCACCATCACTGGCTCTCCGGTCTCCATCGCCCTG
GCCAGTACCTCATCACTGCCTGTCTAGAGACGGCCAAGTCTACCTCTGGGGGACGCC
AGCTCGGCCCCCGCAGACCTGCTGCCCCCTTCTCGCCACCCCTGACGGCCCTGCCACA
GCTCCCCCTGGCCTGCTGGGCACACCCTATGCCATCTCCCTCTCCAACCTCATCGGCCTC
AAGCCCATGCCCTTCTTGGCTTTACCACCTGCTTCCCCAGGGCCGCCCGGGGCTTGGCG
GCCTACACTGCCAAGATGGCAGCAGCTAATGGGAGCAAGAAGGCTGAGCGGCAGAAATTC
TCCCCCTACTGA

```

Restriction Sites:	Please inquire
ACCN:	NM_001174100



[View online >](#)

OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001174100.1</u> , <u>NP_001167571.1</u>
RefSeq Size:	2188 bp
RefSeq ORF:	1212 bp
Locus ID:	57060
UniProt ID:	<u>P57723</u>
Cytogenetics:	3p21.2
Gene Summary:	<p>This gene encodes a member of the KH-domain protein subfamily. Proteins of this subfamily, also referred to as alpha-CPs, bind to RNA with a specificity for C-rich pyrimidine regions. Alpha-CPs play important roles in post-transcriptional activities and have different cellular distributions. This gene is induced by the p53 tumor suppressor, and the encoded protein can suppress cell proliferation by inducing apoptosis and cell cycle arrest in G(2)-M. This gene's protein is found in the cytoplasm, yet it lacks the nuclear localization signals found in other subfamily members. Multiple alternatively spliced transcript variants have been described, but the full-length nature for only some has been determined. [provided by RefSeq, Jul 2008]</p> <p>Transcript Variant: This variant (5) differs in the 5' UTR, and includes an additional in-frame exon in the central coding region, compared to variant 1. The encoded isoform (c) is longer than isoform a. Variants 3, 4 and 5 encode the same isoform.</p>