

## Product datasheet for SC328715

### Repulsive Guidance Molecule A (RGMA) (NM\_001166287) Human Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Repulsive Guidance Molecule A (RGMA) (NM_001166287) Human Untagged Clone
Tag:	Tag Free
Symbol:	RGMA
Synonyms:	RGM
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC328715 representing NM_001166287. Blue=Insert sequence Red=Cloning site Green=Tag(s)


```
GCTCGTTTGTAGTGAACCGTCAGAATTTTGTAAATACGACTACTATAGGGCGCCGGGAATTCGTGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGGGTATGGGGAGAGGGGCAGGACGTTTCAGCCCTGGGATTCTGGCCGACCCCTCGCCTTCTCTCTGC
AGCTTCCCGCAGCCACCTCCCGTGCAAGATCCTCAAGTGAACCTCTGAGTTCTGGAGCGCCACGTCG
GGCAGCCACGCCCCAGCCTCAGACGACACCCCCGAGTTCTGTGCAGCCTTGGCAGCTACGCCCTGTGC
ACGCGGGGACGGCCCGCACCTGCCGGGTGACCTGGCCTACCACTCGGCCGTCATGGCATAGAGGAC
CTCATGAGCCAGCACAAGTCTCAAGGATGGCCCCACCTCGCAGCCAGCCTGCGCAGCTCCCACCG
GCCGGAGACAGCCAGGAGCGCTCGGACAGCCCCGAGATCTGCCATTACGAGAAGAGCTTTCACAAGCAC
TCGGCCACCCCAACTACACGACTGTGGCCTTTCGGGGACCCACACCTCAGGACTTTCACCGACCGC
TTCCAGACCTGCAAGGTGCAGGGCGCCTGGCCGCTCATCGACAATAATTACCTGAACGTGCAGGTCACC
AACACGCCTGTGCTGCCCGCTCAGCGGCCACTGCCACCAGCAAGCTCACCATCATCTTCAAGAACTTC
CAGGAGTGTGGACCAGAAGGTGTACCAGGCTGAGATGGACGAGCTCCCGGCCGCTTCGTGGATGGC
TCTAAGAACGGTGGGACAAGCAGGGGCAACAGCCTGAAGTCACTGAGAAGGTGTGAGGCCAGCAC
GTGGAGATCCAGGCCAAGTACATCGGCACCACCATCGTGGTGCGCCAGGTGGGCCGCTACCTGACCTTT
GCCGTCCGCATGCCAGAGGAAGTGGTCAATGCTGTGGAGGACTGGGACAGCCAGGCTCTACCTCTGC
CTGCGGGGCTGCCCCCTCAACCAGCAGATCGACTTCCAGGCTTCCACACCAATGCTGAGGGCACCGGT
GCCCGCAGGCTGGCAGCCGCGCAGCCTGCACCCACAGCCCCGAGACCTTCCCATACGAGACAGCCGTG
GCCAAGTGAAGGAGAAGCTGCCGGTGGAGGACCTGTAATACCAGGCTGCGTCTTCGACCTCCTCACC
ACGGGCGACGTGAACCTTCACTGCGCCCTACTACGCGTTGGAGGATGTCAAGATGCTCCACTCCAAC
AAAGACAACTGCACCTGTATGAGAGGACTCGGGACCTGCCAGGCAGGGCGGCTGCGGGGCTGCCCTG
GCCCGCGGCCCTCCTGGGCGCCCTCGTCCCGCTCCTGGCCCTGCTCCCTGTGTTCTGCTAG
AGCGGACCGACGCTACGCGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGAT
ATCCTGGATTACAAGGATGACGACGATAAGGTTTAA
```



[View online »](#)

Restriction Sites: SgfI-RsrII

Plasmid Map:



ACCN: NM\_001166287

Insert Size: 1305 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

**OTI Annotation:** This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM\\_001166287.1](#)

RefSeq Size: 3174 bp

RefSeq ORF: 1305 bp

Locus ID: 56963

UniProt ID: [Q96B86](#)

Cytogenetics: 15q26.1

MW: 47.5 kDa

**Gene Summary:** This gene encodes a member of the repulsive guidance molecule family. The encoded protein is a glycosylphosphatidylinositol-anchored glycoprotein that functions as an axon guidance protein in the developing and adult central nervous system. This protein may also function as a tumor suppressor in some cancers. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Oct 2009]

Transcript Variant: This variant (3) differs in the 5' UTR, lacks a portion of the 5' coding region, and initiates translation at a downstream start codon, compared to variant 1. The encoded isoform (2) has a shorter N-terminus, compared to isoform 1. Variants 2, 3, 5 and 6 encode the same isoform (2). Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments. Variants 2 and 3 encode the same isoform (2).