

Product datasheet for **SC328791**

SHMT2 (NM_001166359) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	SHMT2 (NM_001166359) Human Untagged Clone
Tag:	Tag Free
Symbol:	SHMT2
Synonyms:	GLYA; HEL-S-51e; NEDCASB; SHMT
Mammalian Cell Selection:	None
Vector:	<u>pCMV6-XL5</u>
E. coli Selection:	Ampicillin (100 ug/mL)
Fully Sequenced ORF:	>NCBI ORF sequence for NM_001166359, the custom clone sequence may differ by one or more nucleotides

```

ATGGCCATTCGGGCTCAGCACAGCAACGCAGCCCAGACTCAGACTGGGGAAGCAAACAGG
GGCTGGACAGGCCAGGAGAGCCTGTCGGACAGTGATCCTGAGATGTGGGAGTTGCTGCAG
AGGGAGAAGGACAGGCAGTGTCTGGCCTGGAGCTCATTGCCTCAGAGAATTCTGCAGC
CGAGCTGCCTGGAGGCCCTGGGGTCTGTCTGAACAACAAGTACTCGGAGGGTTATCCT
GGCAAGAGATACTATGGGGGAGCAGAGGTGGTGGATGAAATTGAGCTGCTGTGCCAGCGC
CGGGCCTTGAAGCCTTTGACCTGGATCCTGCACAGTGGGGAGTCAATGTCCAGCCCTAC
TCCGGGTCCCAGCCAACCTGGCCGTCTACACAGCCCTTCTGCAACCTCACGACCCGATC
ATGGGGCTGGACCTGCCGATGGGGGCCATCTACCCACGGCTACATGTCTGACGTCAAG
CGGATATCAGCCACGTCCATCTTCTCGAGTCTATGCCCTATAAGCTCAACCCAAAACCT
GGCCTCATTGACTACAACCAGCTGGCACTGACTGCTCGACTTTTCCGGCCACGGCTCATC
ATAGCTGGCACCAGCGCCTATGCTCGCCTCATTGACTACGCCCGCATGAGAGAGGTGTGT
GATGAAGTCAAAGCACACCTGCTGGCAGACATGGCCCACATCAGTGGCCTGGTGGCTGCC
AAGGTGATTCCTCGCCTTTCAAGCACGCGGACATCGTCACCACCACTACTCACAAAGACT
CTTCGAGGGGCCAGGTGAGGGCTCATCTTCTACCGGAAAGGGGTGAAGGCTGTGGACCC
AAGACTGGCCGGGAGATCCCTTACACATTTGAGGACCGAATCAACTTTGCCGTGTCCCA
TCCCTGCAGGGGGGCCCCACAATCATGCCATTGCTGCAGTAGCTGTGGCCCTAAAGCAG
GCCTGCACCCCATGTTCCGGGAGTACTCCCTGCAGTTCTGAAGAATGCTCGGGCCATG
GCAGATGCCCTGCTAGAGCGAGGCTACTACTGTTATCAGGTGGTACTGACAACCACCTG
GTGCTGGTGGACCTGCGGCCAAGGGCCTGGATGGAGCTCGGGCTGAGCGGGTGCTAGAG
CTTGATCCATCACTGCCAACAAGAACCCTGTCTGGAGACCGAAGTGCATCACACCG
GGCGGCTGCGGCTTGGGGCCCCAGCCTTAACTTCTCGACAGTTCGGTGAGGATGACTTC
CGGAGAGTTGTGGACTTTATAGATGAAGGGTCAACATTGGCTTAGAGGTGAAGAGCAAG
ACTGCAAGCTCCAGGATTTCAAATCCTTCTGCTTAAGGACTCAGAAACAAGTCAGCGT
CTGGCCAACCTCAGGCAACGGGTGGAGCAGTTTCCAGGGCCTTCCCATGCCTGGTTTT
GATGAGCATTGA

```



[View online >](#)

Restriction Sites:	Please inquire
ACCN:	NM_001166359
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001166359.1</u> , <u>NP_001159831.1</u>
RefSeq Size:	2149 bp
RefSeq ORF:	1452 bp
Locus ID:	6472
UniProt ID:	<u>P34897</u>
Cytogenetics:	12q13.3
Protein Pathways:	Cyanoamino acid metabolism, Glycine, serine and threonine metabolism, Metabolic pathways, Methane metabolism, One carbon pool by folate
Gene Summary:	This gene encodes the mitochondrial form of a pyridoxal phosphate-dependent enzyme that catalyzes the reversible reaction of serine and tetrahydrofolate to glycine and 5,10-methylene tetrahydrofolate. The encoded product is primarily responsible for glycine synthesis. The activity of the encoded protein has been suggested to be the primary source of intracellular glycine. The gene which encodes the cytosolic form of this enzyme is located on chromosome 17. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Oct 2009] Transcript Variant: This variant (5) differs in the 5' UTR, lacks a portion of the 5' coding region, and initiates translation at a downstream start codon, compared to variant 1. The encoded isoform (3) has a shorter N-terminus, compared to isoform 1. Variants 3, 4, and 5 encode the same isoform 3. Translation efficiency is uncertain from this alternative start codon, and there is rapid turnover of the mitochondrial protein in the cytoplasm prior to mitochondrial import due to its shortened import presequence (PMID 8999870).