

## Product datasheet for **SC328836**

### HCK (NM\_001172132) Human Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	HCK (NM_001172132) Human Untagged Clone
Tag:	Tag Free
Symbol:	HCK
Synonyms:	JTK9; p59Hck; p61Hck
Mammalian Cell Selection:	None
Vector:	<u>pCMV6-XL5</u>
E. coli Selection:	Ampicillin (100 ug/mL)



[View online »](#)

<b>Fully Sequenced ORF:</b>	<p>&gt;NCBI ORF sequence for NM_001172132, the custom clone sequence may differ by one or more nucleotides</p> <pre> ATGATGGGGTGCATGAAGTCCAAGTTCCTCCAGGTCGGAGGCAATACATTCTCAAAAAC GAAACAGCGCCAGCCACACTGTCCTGTGTACGTGCCGATCCACATCCACCATCAAG CCGGGGCTAATAGCCACAACAGCAACACACAGGAATCAGGGAGGCAGGCTCTGAGGAC ATCATCGTGGTTGCCCTGTATGATTACGAGGCCATTACCACGAAGACCTCAGCTTCCAG AAGGGGACCAGATGGTGGTCTAGAGGAATCCGGGAGTGGTGGAAAGGCTCGATCCCTG GCCACCCGGAAGGAGGGCTACATCCCAAGCAACTATGTCGCCCGCGTTGACTCTCTGGAG ACAGAGGAGTGGTTTTTCAAGGGCATCAGCCGGAAGGACGAGAGCGCCAACCTGTGGCT CCCGGCAACATGCTGGGCTCCTTCATGATCCGGGATAGCGAGACCACTAAAGGAAGCTAC TCTTTGTCGTCGAGACTACGACCTCGGCAGGGAGATACCGTGAAACATTACAAGATC CGGACCTGGACAACGGGGGCTTCTACATATCCCCCGAAGCACCTTCAGCACTCTGCAG GAGCTGGTGGACCACTACAAGAAGGGGAACGACGGGCTCTGCCAGAACTGTCGGTGCC TGCATGTCTTCAAGCCCCAGAAGCCTTGGGAGAAAGATGCCTGGGAGATCCCTCGGGAA TCCCTCAAGCTGGAGAAGAACTTGGAGCTGGGCAGTTTGGGGAAGTCTGGATGGCCACC TACAACAAGCACCAAGGTGGCAGTGAAGACGATGAAGCCAGGAGCATGTCGGTGGAG GCCTTCCTGGCAGAGGCCAACGTGATGAAAACCTGTCAGCATGACAAGCTGGTCAAACCT CATGCGGTGGTCACCAAGGAGCCCATCTACATCATCAGGAGTTCATGGCCAAAGGAAGC TTGCTGGACTTTCTGAAAAGTGATGAGGGCAGCAAGCAGCCATTGCCAAAACCTATTGAC TTCTCAGCCAGATTGCAGAAGGCATGGCCTTCATCGAGCAGAGGAACATACCCACCGA GACCTCCGAGCTGCCAACATCTTGGTCTCTGCATCCCTGGTGTGAAGATTGTGACTTT GGCCTGGCCCGGGTCATTGAGGACAACGAGTACACGGCTCGGGAAGGGGCAAGTTCCCC ATCAAGTGGACAGCTCCTGAAGCCATCAACTTTGGCTCCTTCACCATCAAGTCAGACGTC TGGTCCTTTGGTATCCTGCTGATGGAGATCGTCACCTACGGCCGGATCCCTTACCCAGGG ATGTCAAACCCTGAAGTGATCCGAGCTCTGGAGCGTGGATACCGGATGCCTCGCCAGAG AACTGCCAGAGGAGCTCTACAACATCATGATGCGCTGCTGAAAAACCGTCCGGAGGAG CGGCCGACCTTGAATACATCCAGAGTGTGCTGGATGACTTCTACACGGCCACAGAGAGC CAGTACCAACAGCAGCCATGA </pre>
<b>Restriction Sites:</b>	Please inquire
<b>ACCN:</b>	NM_001172132
<b>OTI Disclaimer:</b>	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
<b>OTI Annotation:</b>	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

<b>Reconstitution Method:</b>	<ol style="list-style-type: none"> <li>1. Centrifuge at 5,000xg for 5min.</li> <li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>3. Close the tube and incubate for 10 minutes at room temperature.</li> <li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
<b>RefSeq:</b>	<u>NM_001172132.1</u> , <u>NP_001165603.1</u>
<b>RefSeq Size:</b>	2234 bp
<b>RefSeq ORF:</b>	1521 bp
<b>Locus ID:</b>	3055
<b>Cytogenetics:</b>	20q11.21
<b>Protein Families:</b>	Druggable Genome, Protein Kinase
<b>Protein Pathways:</b>	Chemokine signaling pathway, Fc gamma R-mediated phagocytosis
<b>Gene Summary:</b>	<p>The protein encoded by this gene is a member of the Src family of tyrosine kinases. This protein is primarily hemopoietic, particularly in cells of the myeloid and B-lymphoid lineages. It may help couple the Fc receptor to the activation of the respiratory burst. In addition, it may play a role in neutrophil migration and in the degranulation of neutrophils. Multiple isoforms with different subcellular distributions are produced due to both alternative splicing and the use of alternative translation initiation codons, including a non-AUG (CUG) codon. [provided by RefSeq, Feb 2010]</p> <p>Transcript Variant: This variant (3) differs in its 5' UTR and uses an alternate AUG translation start codon, compared to variant 1. The encoded isoform (e) has a distinct and shorter N-terminus, compared to isoform a.</p>