

Product datasheet for SC329409

p53 (TP53) (NM_001126118) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: p53 (TP53) (NM_001126118) Human Untagged Clone

Tag: Tag Free Symbol: TP53

Synonyms: BCC7; BMFS5; LFS1; P53; TRP53

Vector: pCMV6-Entry (PS100001)

Fully Sequenced ORF: >SC329409 representing NM_001126118.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

TTCAAGACAGAAGGGCCTGACTCAGACTGA

Restriction Sites: Sgfl-Mlul




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Plasmid Map:



ACCN: NM_001126118

Insert Size: 1065 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001126118.1</u>

RefSeq Size: 2708 bp



p53 (TP53) (NM_001126118) Human Untagged Clone - SC329409

 RefSeq ORF:
 1065 bp

 Locus ID:
 7157

 UniProt ID:
 P04637

 Cytogenetics:
 17p13.1

Protein Families: Druggable Genome, Stem cell - Pluripotency, Transcription Factors

Protein Pathways: Amyotrophic lateral sclerosis (ALS), Apoptosis, Basal cell carcinoma, Bladder cancer, Cell cycle,

Chronic myeloid leukemia, Colorectal cancer, Endometrial cancer, Glioma, Huntington's disease, MAPK signaling pathway, Melanoma, Neurotrophin signaling pathway, Non-small cell lung cancer, p53 signaling pathway, Pancreatic cancer, Pathways in cancer, Prostate cancer,

Small cell lung cancer, Thyroid cancer, Wnt signaling pathway

MW: 39.3 kDa

Gene Summary: This gene encodes a tumor suppressor protein containing transcriptional activation, DNA

binding, and oligomerization domains. The encoded protein responds to diverse cellular stresses to regulate expression of target genes, thereby inducing cell cycle arrest, apoptosis, senescence, DNA repair, or changes in metabolism. Mutations in this gene are associated with a variety of human cancers, including hereditary cancers such as Li-Fraumeni syndrome.

Alternative splicing of this gene and the use of alternate promoters result in multiple

transcript variants and isoforms. Additional isoforms have also been shown to result from the use of alternate translation initiation codons from identical transcript variants (PMIDs:

12032546, 20937277). [provided by RefSeq, Dec 2016]

Transcript Variant: This variant (8, also known as p53l2) differs in the 5' UTR, and uses an inframe downstream start codon, compared to variant 1. The encoded isoform (g, also known as delta40p53alpha or deltaNp53) has a shorter N-terminus, compared to isoform a. This

variant is supported by data in PMID:21112961.