

Product datasheet for SC329419

RPH3AL (NM 001190412) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: RPH3AL (NM_001190412) Human Untagged Clone

Tag: Tag Free
Symbol: RPH3AL
Synonyms: NOC2

Vector: pCMV6-Entry (PS100001)

Fully Sequenced ORF: >SC329419 representing NM_001190412.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

GCTCCAGCAGGCCCCTCCAGCTGCCTGGGCTGA

Restriction Sites: Sgfl-Mlul


OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Plasmid Map:



ACCN: NM_001190412

Insert Size: 861 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.



RPH3AL (NM_001190412) Human Untagged Clone - SC329419

RefSeq: <u>NM 001190412.1</u>

RefSeq Size: 2667 bp
RefSeq ORF: 861 bp
Locus ID: 9501
UniProt ID: Q9UNE2
Cytogenetics: 17p13.3

Protein Families: Secreted Protein

MW: 31.2 kDa

Gene Summary: The protein encoded by this gene plays a direct regulatory role in calcium-ion-dependent

exocytosis in both endocrine and exocrine cells and plays a key role in insulin secretion by pancreatic cells. This gene is likely a tumor suppressor. Alternative splicing results in multiple

transcript variants encoding distinct isoforms. [provided by RefSeq, Jun 2010]

Transcript Variant: This variant (3) lacks an in-frame exon in the coding region, compared to variant 1, and encodes a shorter isoform (2) than isoform 1. Variants 3 and 4 encode the

same isoform (2).