

Product datasheet for **SC329574**

DAP3 (NM_001199850) Human Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: DAP3 (NM_001199850) Human Untagged Clone
Tag: Tag Free
Symbol: DAP3
Synonyms: bMRP-10; DAP-3; MRP-S29; MRPS29; S29mt
Vector: pCMV6-Entry (PS100001)
Fully Sequenced ORF: >SC329574 representing NM_001199850.
 Blue=Insert sequence Red=Cloning site Green=Tag(s)


```

ATGATGCTGAAAGGAATAACAAGGCTTATCTCTAGGATCCATAAGTTGGACCCTGGGCGTTTTTACAC
ATGGGGACCCAGGCTCGCCAAAGCATTGCTGCTCACCTAGATAACCAGGTTCCAGTTGAGAGTCCGAGA
GCTATTTCCCGCACCAATGAGAATGACCCGGTGAAGACATTCAGTGAAGCTTGCCTGATGGTAAGGAAA
CCAGCCCTAGAATTCTGCATTACCTGAAAAACACCAGTTTTGCTTATCCAGCTATACGATATCTTCTG
TATGGAGAGAAGGAACAGGAAAAACCCTAAGTCTTTGCCATGTTATTCAATTTCTGTGCAAAACAGGAC
TGGCTGATACTACATATCCAGATGCTCATCTTTGGGTGAAAAATGTGCGGGATCTTCTGCAGTCCAGC
TACAACAAACAGCGCTTTGATCAACCTTTAGAGGCTTCAACCTGGCTGAAGAATTTCAAACACTACAAT
GAGCGCTTCTGAACCAGATAAAAGTTCAAGAGAAGTATGTCTGGAATAAGAGAGAAAGCACTGAGAAA
GGGAGTCTCTGGGAGAAGTGGTTGAACAGGGCATAACACGGGTGAGGAACGCCACAGATGCAGTTGGA
ATTGTGCTGAAAGAGCTAAAGAGGCAAAGTTCTTTGGGTATGTTTCACCTCCTAGTGGCCGTGGATGGA
ATCAATGCTCTTTGGGGAAGAACCCTCTGAAAAGAGAAGATAAAAGCCCGATTGCCCCGAGGAATTA
GCACTTGTTCACAACCTTGAGGAAAATGATGAAAATGATTGGCATGGAGGCGCCATTGTGTGGCTTTG
AGCCAGACTGGTCTCTCTTTAAGCCCCGAAAGCCTATCTGCCCCAGGAGTTGCTGGGAAAGGAAGGA
TTTGATGCCCTGGATCCCTTTATCCCATCTGGTTTCCAATAAACCAGGAATTTGAAAGTTGT
ATTCAGTATTATTTGAAAACAATTGGCTTCAACATGAGAAAGCTCTACAGAAGAAGGGAAAAAGAG
CTGCTGTTCTAAGTAACGCGAACCCCTCGCTGCTGGAGCGGCACTGTGCCTACCTCAA
  
```

Restriction Sites: SgfI-MluI



[View online »](#)

Plasmid Map:


ACCN: NM_001199850

Insert Size: 1095 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001199850.1](#)

RefSeq Size: 1966 bp

RefSeq ORF: 1095 bp

Locus ID: 7818

UniProt ID: [P51398](#)

Cytogenetics: 1q22

Protein Families: Druggable Genome

MW: 41.7 kDa

Gene Summary: Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a 28S subunit protein that also participates in apoptotic pathways which are initiated by tumor necrosis factor-alpha, Fas ligand, and gamma interferon. This protein potentially binds ATP/GTP and might be a functional partner of the mitoribosomal protein S27. Multiple alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. Pseudogenes corresponding to this gene are found on chromosomes 1q and 2q. [provided by RefSeq, Dec 2010]

Transcript Variant: This variant (4) lacks an in-frame exon in the CDS, as compared to variant 2. The resulting isoform (2) lacks an internal segment, as compared to isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.