

Product datasheet for SC329618

ROC2 (RNF7) (NM_001201370) Human Untagged Clone

Product data:


Product Type:	Expression Plasmids
Product Name:	ROC2 (RNF7) (NM_001201370) Human Untagged Clone
Tag:	Tag Free
Symbol:	RNF7
Synonyms:	CKBBP1; rbx2; ROC2; SAG
Vector:	pCMV6-Entry (PS100001)
Fully Sequenced ORF:	>SC329618 representing NM_001201370. Blue=Insert sequence Red=Cloning site Green=Tag(s)

```

ATGGCCGACGTGGAAGACGGAGAGGAAACCTGCGCCCTGGCCTCTCACTCCGGGAGCTCAGGCTCCAAG
TCGGGAGCGACAAGATGTTCTCCCTCAAGAAGTGAACGCGGTGGCCATGTGGAGCTGGGACGTGGAG
TGCGATACGTGCCCATCTGCAGGGTCCAGGTGATGGTGGTCTGGGGAGAATGTAATCATTCTCCAC
AACTGCTGCATGCCCTGTGGGTGAAACAGAACAATCGCTGCCCTCTCTGCCAGCAGGACTGGGTGGTC
CAAAGAATCGGCAAAATGA
  
```

Restriction Sites: SgfI-MluI

Plasmid Map:



ACCN: NM_001201370

Insert Size: 294 bp



[View online >](#)

OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001201370.1</u>
RefSeq Size:	1962 bp
RefSeq ORF:	294 bp
Locus ID:	9616
UniProt ID:	<u>Q9UBF6</u>
Cytogenetics:	3q23
Protein Families:	Druggable Genome
Protein Pathways:	Ubiquitin mediated proteolysis
MW:	10.9 kDa
Gene Summary:	<p>The protein encoded by this gene is a highly conserved ring finger protein. It is an essential subunit of SKP1-cullin/CDC53-F box protein ubiquitin ligases, which are a part of the protein degradation machinery important for cell cycle progression and signal transduction. This protein interacts with, and is a substrate of, casein kinase II (CSNK2A1/CKII). The phosphorylation of this protein by CSNK2A1 has been shown to promote the degradation of IκBα (CHUK/IKK-α/IKBKA) and p27Kip1(CDKN1B). Alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]</p> <p>Transcript Variant: This variant (4) lacks an alternate in-frame exon compared to variant 1. The resulting isoform (4) has the same N- and C-termini but is shorter compared to isoform 1.</p> <p>Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.</p>