

Product datasheet for SC329630

Proprotein Convertase 2 (PCSK2) (NM_001201529) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Proprotein Convertase 2 (PCSK2) (NM_001201529) Human Untagged Clone
Tag:	Tag Free
Symbol:	PCSK2
Synonyms:	NEC-2; NEC 2; NEC2; PC2; SPC2
Vector:	pCMV6-Entry (PS100001)
Fully Sequenced ORF:	>SC329630 representing NM_001201529. Blue=Insert sequence Red=Cloning site Green=Tag(s)

```

ATGAAGGGTGGTTGTGTCTCCAGTGGAAAGCGGCCGCCGGTTCCCTCTTCTGTGTATGGTTTTTGC
TCTGCTGAGCGACCGGTCTTACGAATCATTTTTCTGTGGAGTTGCATAAAGGGGAGAGACAAAGCT
CGCCAAGTTGCAGCAGAACACGGCTTTGGAGTCCGAAAGGTAAGATGGCTTTCAGCAGGAAGGATTT
GACCGAAAAAGCGAGGTTACAGAGACATCAATGAGATCGACATCAACATGAACGATCCTCTTTTTACA
AAGCAGTGGTATCTGATCAATACTGGGCAAGCTGATGGCACTCCTGGCCTTGATTTGAATGTGGTGAA
GCCTGGGAGCTGGGATACACAGGAAAGGTGTTACCATTGGAATTATGGATGATGGGATTGACTATCTC
CACCCGGACCTGGCCTCCAATAATGCCGAAGCAAGTTACGACTTCAGCAGCAACGACCCCTATCCT
TACCCTCGGTACACAGATGACTGGTTAACAGCCACGGGACCCGATGTGCAGGAGAAGTTTCTGCTGCC
GCCAACAAACAATATCTGTGGAGTTGGAGTAGCATACAACCTCCAAGGTTGCAGGCATCCGGATGCTGGAC
CAGCCATTCATGACAGACATCATCGAGGCTCCTCCATCAGTCATATGCCACAGCTGATTGACATCTAC
AGCGCCAGCTGGGGCCCCACAGACAACGGCAAGACAGTGGATGGGCCCGGGAGCTCACGCTGCAGGCC
ATGGCCGATGGCGTGAACAAGGGCCGCGCGGCAAGGCAGCATCTACGTGTGGCCTCCGGGGACGGC
GGCAGCTATGACGACTGCAACTGCGACGGCTACGCCTCCAGCATGTGGACCATCTCCATCAACTCAGCC
ATCAACGACGGCAGGACTGCCCTGTACGACGAGAGCTGCTTTCCACCTTGCTTCCACCTTCAGCAAC
GGGAGGAAAAGGAACCCCGAGGCCGGTGTGGCAACCACAGATTTGTACGGCAACTGCACTCTGAGGCAT
TCTGGGACATCTGCAGCTGCCCCGAGGACGCTGGTGTGTTTGCAGTGGCTCTGGAGGCTAACCTGGGT
CTGACCTGGCGGGACATGCAGCATCTGACTGTGCTCACCTCCAAACGGAACCAGCTTCACGACGAGGTC
CATCAGTGGCGGCGCAATGGGGTCGGCCTGGAATTTAATCACCTCTTTGGCTACGGGGTCTTTGATGCA
GGTGCCATGGTAAAAATGGCTAAAGACTGGAAAACCGTGCCTGAGAGATTCCACTGTGTGGGAGGCTCC
GTGCAGGACCCTGAGAAAATACCATCCACTGGCAAGTTGGTGTGACACTCACAACCGACGCTGTGAG
GGGAAGGAAAATTTGTCCGCTACCTGGAGCATGTCCAGGCTGTATCACGGTCAACGCAACCAGAAGA
GGAGACCTGAACATCAACATGACTTCCCCTATGGGCACCAAGTCCATTTTGTGAGCCGGCTCCAAGG
GATGACGACTCCAAGGTGGGCTTTGACAAGTGGCCTTTTATGACCACTCACACGTGGGGGAAGACGCC
CGAGGCACCTGGACCCTGGAGCTGGGATTTGTCCGACGCCCCCGAGAAGGGGGTGTGTAAGGAGTGG
ACCCTGATGCTGCATGGCACTCAGAGTGGCCCGTACATCGACCAGGTGGTGCAGGATTACAGTCCAAG
TTGGCCATGTCCAAGAAAGAGGAGCTGGAGGAAGAGCTGGACGAAGCCGTGGAGAGAAGCCTGAAAAGC
ATCCTTAAACAAGAACTAG


```



[View online »](#)

Restriction Sites: SgfI-MluI

Plasmid Map:



ACCN: NM_001201529

Insert Size: 1812 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001201529.1](#)

RefSeq Size: 4682 bp

RefSeq ORF: 1812 bp

Locus ID: 5126

UniProt ID: [P16519](#)

Cytogenetics: 20p12.1

Protein Families: Druggable Genome, Protease, Secreted Protein

MW: 66.3 kDa

Gene Summary:

This gene encodes a member of the subtilisin-like proprotein convertase family, which includes proteases that process protein and peptide precursors trafficking through regulated or constitutive branches of the secretory pathway. The protein undergoes an initial autocatalytic processing event and interacts with a neuroendocrine secretory protein in the ER, exits the ER and sorts to secretory granules, where it is cleaved and catalytically activated during intracellular transport. The encoded protease is packaged into and activated in dense core secretory granules and expressed in the neuroendocrine system and brain. This gene encodes one of the seven basic amino acid-specific members which cleave their substrates at single or paired basic residues. It functions in the proteolytic activation of polypeptide hormones and neuropeptides precursors. Single nucleotide polymorphisms in this gene may increase susceptibility to myocardial infarction and type 2 diabetes. This gene may also play a role in tumor development and progression. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Jan 2014]

Transcript Variant: This variant (2) lacks an alternate in-frame exon in the 5' coding region, compared to variant 1. The encoded isoform (2) is shorter than isoform 1. **Sequence Note:** This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.